


Journal Pre-proof

The influence of dietary crude protein concentrations, grain types and arginine:lysine ratios on the performance of broiler chickens

Shemil P. Macelline, Peter V. Chrystal, Chanon Inanan, Mehdi Toghyani, Peter H. Selle, Sonia Yun Liu



PII: S2405-6545(23)00065-3

DOI: <https://doi.org/10.1016/j.aninu.2023.05.007>

Reference: ANINU 747

To appear in: *Animal Nutrition Journal*

Received Date: 18 August 2022

Revised Date: 11 May 2023

Accepted Date: 22 May 2023

Please cite this article as: Macelline SP, Chrystal PV, Inanan C, Toghyani M, Selle PH, Liu SY, The influence of dietary crude protein concentrations, grain types and arginine:lysine ratios on the performance of broiler chickens, *Animal Nutrition Journal*, <https://doi.org/10.1016/j.aninu.2023.05.007>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 The Authors. Publishing services by Elsevier B.V. on behalf of KeAi Communications Co. Ltd.

Author contributions

All six authors contributed towards the completion of this study and have read and approved this manuscript. **Sonia Yun Liu** was the principal investigator of the relevant project and is the corresponding author. **Peter Vincent Chrystal** formulated the diets. **Shemil Priyan Macelline** and **Mehdi Toghyani** conducted and supervised the feeding study. **Peter Henry Selle** and **Shemil Priyan Macelline** completed the statistical analyses. **Chanon Inanan** completed the initial manuscript, which was completed by **Peter Henry Selle** and **Shemil Priyan Macelline** and **Sonia Yun Liu** was responsible for the final editing and submission of the manuscript.

1 **The influence of dietary crude protein concentrations, grain types and arginine:lysine ratios on**
2 **the performance of broiler chickens**

3
4 Shemil P. Macelline^{a,b}, Peter V. Chrystal^{a,b,c}, Chanon Inanan^a, Mehdi Toghyani^{a,b}, Peter H. Selle^{b,d},
5 Sonia Yun Liu^{a,b,*}

6
7 ^aSchool of Life and Environmental Sciences, Faculty of Science, The University of Sydney.
8 Camperdown 2006 NSW, Australia.

9 ^bPoultry Research Foundation within The University of Sydney. Camden 2570 NSW, Australia.

10 ^cComplete Feed Solutions, Howick 2145, New Zealand.

11 ^dSydney School of Veterinary Science, Faculty of Science, The University of Sydney. Camden 2570
12 NSW, Australia.

13
14 *Corresponding author.

15 Email address: sonia.liu@sydney.edu.au (S.Y. Liu)

16

17 Abstract

18 The objective of this study was to investigate the effects of dietary crude protein (CP) concentrations,
19 grain types and arginine:lysine ratios on performance parameters of broiler chickens. The $2 \times 2 \times 2$
20 factorial array of dietary treatments harnessed two CP concentrations (210 and 170 g/kg), two feed
21 grains (wheat and sorghum), and 2 arginine:lysine ratios (104 and 110). Each dietary treatment was
22 offered to 7 replicates of 14 birds per floor pen, a total of 784 off-sex male, Ross 308 broilers, from 14
23 to 35 d post-hatch. The dietary CP reduction compromised weight gain by 10.0% (2078 versus 2310
24 g/bird) as a main effect and FCR by 7.51% (1.474 versus 1.371), subject to an interaction. In a three-
25 way interaction ($P = 0.008$), expanded arginine:lysine ratios improved FCR by 2.30% in 170 g/kg CP,
26 sorghum-based diets but compromised FCR by 2.12% in corresponding wheat-based diets. Sorghum
27 was the more suitable feed grain in reduced-CP diets as sorghum generated significant advantages in
28 weight gain of 7.59% (2154 versus 2002 g/kg) and FCR of 6.94% (1.421 versus 1.527) in birds offered
29 170 g/kg CP diets. Both dietary CP and feed grain generated significant and divergent impacts in
30 apparent ileal digestibility coefficients for the majority of 16 assessed amino acids. Dietary CP
31 reductions increased non-bound amino acid inclusions (NBAA) in wheat-based diets (48.96 versus
32 9.80 g/kg) to a greater extent than sorghum-based diets (35.3 versus 9.50 g/kg) and increasing dietary
33 NBAA inclusions were linearly associated with compromised weight gain ($r = -0.834$; $P < 0.001$) and
34 FCR ($r = 0.862$; $P < 0.001$). Increasing ratios of free arginine to lysine plasma concentrations were
35 linearly ($r = -0.466$; $P = 0.004$) related to improvements in FCR. The implications of the observed
36 outcomes are discussed and possible explanations are advanced.

37

38 **Key words:** Amino acid; Broiler chickens; Energy; Protein; Sorghum; Wheat

39

40 1. Introduction

41 The growing global demand for chicken-meat makes sustainable nutrient utilisation an imperative to
42 guarantee future food security. The global per capita chicken-meat consumption was 14.9 kg in 2022,
43 which represents approximately 44% of total meat consumption, but is projected to increase to 15.1 kg
44 in 2029 (OECD, 2022). Dietary inputs of 471 g of crude protein (CP) and 30.78 MJ metabolisable
45 energy (ME) are required to generate 1 kg of edible chicken-meat, based on performance objectives
46 and nutrient specifications from one breeding company. Therefore, meeting dietary CP (Liu et al.,
47 2021) and energy density (Gopinger et al., 2017) requirements with precision will enhance sustainable
48 chicken-meat production.

49 Wheat and sorghum are the principal feed grains in Australian broiler diets with wheat being
50 dominant. The feasibility of dietary CP reductions from 222 to 193 and 165 g/kg in either maize- or
51 wheat-based diets for broiler chickens was compared in Chrystal et al. (2021) and maize was the more
52 suitable feed grain in reduced-CP diets. The growth performance of birds offered 165 g/kg CP, wheat-
53 based diets was seriously compromised, but the 193 g/kg CP wheat-based diets supported statistically
54 comparable growth performance to the 222 g/kg CP diets. This suggests that CP could be reduced in
55 wheat-based diets by approximately 30 g/kg without negatively influencing growth performance. Birds
56 offered reduced-CP, wheat-based diets performed satisfactorily in Yin et al. (2020), but not in
57 Greenhalgh et al. (2020) and Chrystal et al. (2021). Consequently, the shortfalls of wheat have been
58 reviewed given its apparent inferiority to maize in the context of reduced-CP diets (Selle et al., 2022a).
59 Wheat has higher protein content than maize and requires higher inclusions of non-bound (synthetic,
60 crystalline) amino acids to meet specifications in reduced-CP diets. Wheat has a more rapid starch
61 digestion rate than maize (Giuberti et al., 2012) and contains more soluble non-starch polysaccharides
62 (Bach Knudsen, 1997). It is then relevant that sorghum is similar to maize in these respects and this
63 also applies to amino acid profiles (Selle et al., 2022a). Thus, sorghum may be a more suitable feed

64 grain than wheat in the context of reduced-CP diets and for this reason the two feed grains are compared
65 in the present study.

66 Arginine is an essential amino acid in broiler diets and is involved in multiple physiological
67 pathways in poultry (Castro and Kim, 2020). Unlike most animal species, arginine is essential in
68 poultry and broiler chickens have a comparatively high dietary arginine requirement of 11.0 to 12.5
69 g/kg (Ball et al., 2007). The high requirement stems from high rates of protein deposition in chickens,
70 the lack of endogenous synthesis, and metabolic interactions between arginine and lysine (Ball et al.,
71 2007). An undesirable property of reduced-CP diets is increased lipid deposition as monitored by
72 relative abdominal fat-pad weights. Over a series of three similar studies (Chrystal et al., 2020a,b,c)
73 dietary CP reductions from 202 to 161 g/kg in maize-based diets increased relative fat-pad weights by
74 an average of 71.4% (13.44 versus 7.84 g/kg). However, dietary arginine supplementation has been
75 shown to decrease abdominal fat content in broiler chickens. Fouad et al. (2013) reported that the
76 inclusion of 2.5 g/kg arginine in a maize-soy diet, containing 12.5 g/kg arginine and 11.1 g/kg lysine,
77 significantly decreased fat-pad weights by 20.7% (15.7 versus 19.8 g/kg). Antagonistic arginine-lysine
78 interactions are established in poultry (Austic and Scott, 1975); therefore, the dietary arginine to lysine
79 ratio assumes importance. A dietary arginine to lysine ratio of 108 has been recommended (Wu, 2014),
80 but there are indications that higher ratios may be advantageous (Zampiga et al., 2018; Castro et al.,
81 2020; Corzo et al., 2021). Therefore, dietary arginine to lysine ratios of 104 and 110 were compared
82 in the present study.

83 The present study was designed to compare wheat and sorghum as the feed grain basis of
84 standard- and reduced-CP diets with two arginine:lysine ratios. Thus, the hypotheses tested were that
85 sorghum is a more suitable feed grain than wheat in reduced-CP diets and that elevated arginine:lysine
86 ratios will improve performance of broiler chickens offered reduced-CP diets.

87 2. Material and methods

88 **2.1 Animal ethics statement**

89 This feeding study fully complied with the specific guidelines (2019/1651) approved by the Research
90 Integrity and Ethics Administration of The University of Sydney.

91 **2.2 Diet preparation**

92 An outline of the dietary treatments is included in Table 1. The formulations of the experimental diets
93 were based on near-infrared spectroscopy (NIR) of wheat, sorghum and soybean meal using the
94 AMINOIR Advanced program (Evonik Nutrition & Care GmbH, Hanau, Germany). Sorghum and
95 wheat were mediumly ground (4.0 mm hammer-mill screen) prior to being blended into complete diets
96 which were steam-pelleted through a Palmer PP330 pellet press (Palmer Milling Engineering, Griffith,
97 NSW, Australia) at a conditioning temperature of 80 °C with a conditioner residence time of 14
98 seconds and were then cooled. The composition and nutrient specifications of the experimental diets
99 are shown in Tables 2 and 3, respectively. All diets contained exogenous phytase (Aextra PHY, Danisco
100 Animal Nutrition) and xylanase (Aextra XB, Danisco Animal Nutrition) and 20 g/kg Celite (Celite
101 Corporation, Lompoc, CA, USA) as an inert dietary marker. All diets were formulated to 11.0 g/kg
102 digestible lysine, 13.1 g/kg glycine equivalents and DEB was maintained at 210 mEq/kg. The analysed
103 starch, protein (N) and amino acid concentrations in the 8 dietary treatments are shown in Table 4.
104 There are some discrepancies in the analysed amino acid concentrations, which, as mentioned below,
105 were taken into account to calculate amino acid digestibility coefficients.

106 **2.3 Bird management**

107 A total of 784 male Ross 308 one-day-old chicks were procured from a commercial hatchery and were
108 initially offered a conventional starter diet (230 g/kg CP) from 1 to 13 d post-hatch. At d 14, birds were
109 weighed and distributed to 56 floor-pens to ensure an even body-weight distribution (average body-
110 weight 491 ± 16.0 g/bird per pen). Each of the eight dietary treatments was offered to seven replicate
111 pens (14 birds per pen) from 14 to 35 d post-hatch. The floor pen dimensions were 1.5 m in width and

112 depth. Birds had unrestricted access to feed and water in an environmentally controlled facility under
113 a lighting schedule of 18-h-on and 6-h-off. An initial room temperature of 32 °C was maintained for
114 the first week, which was gradually decreased to 22 °C and kept constant to the end of the experiment.

115 ***2.4 Data and sample collection, chemical analyses, calculations***

116 Growth performance (weight gain, feed intake, FCR) was determined from 14 to 35 d post-hatch.
117 Birds were weighed at d 14 and 35 and feed intakes were monitored over this interval, bodyweights of
118 any dead or culled birds were recorded daily to correct feed intakes on a per pen basis and adjust FCR
119 calculations.

120 At 34 d post-hatch, blood samples were taken from the brachial vein of three representative
121 birds per pen to determine free amino acid concentrations in systemic plasma. Blood samples were
122 centrifuged and decanted plasma samples were held at –80 °C prior to analysis. Concentrations of
123 twenty proteinogenic amino acids were determined using precolumn derivatisation amino acid analysis
124 with 6-aminoquinolyl-*N*-hydroxysuccinimidyl carbamate (AQC; Waters AccQTag Ultra; Waters
125 Australia PL; www.waters.com) followed by separation of the derivatives and quantification by
126 reversed phase ultra-performance liquid chromatography (RP-UPLC). All amino acids were detected
127 by UV absorbance and this procedure is fully described in Selle et al. (2016).

128 At 35 d post-hatch, birds were euthanised by an intravenous injection of sodium
129 pentobarbitone, abdominal cavities opened, and abdominal fat-pads dissected out and weighed and
130 recorded against final body weights. The small intestine was removed and digesta was gently expressed
131 manually from the distal half of the ileum and pooled by cage, homogenised, freeze dried and weighed
132 to determine the apparent digestibility coefficients of starch, crude protein (N) and amino acids. Starch
133 concentrations in diets and digesta samples were determined by using total starch assay kits
134 (Megazyme, Wicklow, Ireland) as described in Mahasukhonthachat et al. (2010) and protein (N)
135 concentrations were determined by methods described in Siriwan et al. (1993). Amino acid

136 concentrations of diets and digesta were determined via 24 h liquid hydrolysis at 110 °C in 6 mol/L
137 HCl followed by analysis of 16 amino acids using the Walters AccQTag Ultra chemistry on a Waters
138 Acquity UPLC. Amino acid analyses were completed as outlined by Cohen and Michaud (1993).
139 Apparent crude protein, starch and amino acid digestibility coefficients in distal jejunum and distal
140 ileum were calculated by the following equation:

$$141 \quad \text{Digestibility coefficient} = \frac{(\text{Nutrient/AIA})_{\text{Diet}} - (\text{Nutrient/AIA})_{\text{Digesta}}}{(\text{Nutrient/AIA})_{\text{Diet}}}$$

142 Some discrepancies in analysed amino acid concentrations were detected, which mainly involved
143 amino acids with high dietary inclusions as non-bound entities. Seven amino acids were included
144 across all diets as both protein-bound and non-bound entities. Significant linear relationships between
145 dietary non-bound amino acid inclusions and analysed concentrations of six amino acids were detected.
146 These included four negative (isoleucine, lysine, threonine, valine) and two positive (methionine,
147 glycine) relationships. These anomalies indicate that non-bound and protein-bound amino acids are
148 not being extracted at identical rates during the analytical procedures. Therefore, calculations of
149 apparent amino acid digestibility coefficients were adjusted by substituting total specified
150 concentrations for the eleven amino acids that were included in experimental diets as non-bound
151 entities for the analysed concentrations. Disappearance rates (g/bird per day) of protein (N), and starch
152 in the distal ileum were calculated as the product of dietary concentrations of nutrient (g/kg), daily
153 feed intake (g/day) from 14 to 35 d post-hatch and the relevant digestibility coefficient. Carcass yields
154 were obtained from the manual processing of four birds selected at random from each pen. Breast and
155 leg quarters were removed in their entirety, weighed and recorded against final body weights.

156 *2.5 Statistical analysis*

157 Experimental data were analysed as $2 \times 2 \times 2$ factorial array by analyses of variance using the JMP
158 Pro 16.0 software package (SAS Institute Inc. JMP Software. Cary, NC). The model used for the
159 analyses of variance was as follows:

$$Y_{ijkl} = \mu + \alpha_i + \beta_j + \gamma_k + (\alpha\beta)_{ij} + (\alpha\gamma)_{ik} + (\beta\gamma)_{jk} + (\alpha\beta\gamma)_{ijk} + \varepsilon_{ijkl}$$

161 Where;

162 Y : t -th response observed for treatment i, j, k ; μ : overall mean; α_i : effect on the response due to the
 163 i th level of factor 1; β_j : effect on the response due to the j th level of factor 2; γ_k : effect on the response
 164 due to the k th level of factor 3; ε_{ijkl} : independent random variables

165 Two-way interactions: $\alpha\beta, \alpha\gamma, \beta\gamma$; Three-way interaction: $\alpha\beta\gamma$

166 Linear and quadratic regressions and Pearson correlations were established when considered
 167 appropriate. Pen means were the experimental units and a probability level of less than 5% was
 168 considered statistically significant.

169 3. Results

170 The effects of dietary treatments on growth performance and relative abdominal fat pad-weights shown
 171 as Table 5. A feed grain by CP interaction ($P = 0.013$) was observed for weight gain because birds
 172 offered 170 g/kg sorghum-based diets had a 7.59% advantage (2154 versus 2002 g/bird) over their
 173 wheat-based counterparts. Reducing dietary CP depressed feed intake by 3.50% (3057 versus 3168
 174 g/bird; $P < 0.001$). There was a three-way treatment interaction ($P = 0.008$) for FCR, where there were
 175 no statistical differences between birds offered 210 g/kg CP diets. However, sorghum-based diets
 176 supported an FCR of 1.421 in comparison to 1.527 for wheat-based diets in 170 g/kg CP diets.
 177 Moreover, expanded arginine:lysine ratios significantly improved FCR with sorghum-based diets by
 178 2.30% (1.404 versus 1.437) but compromised FCR by 2.12% (1.543 versus 1.511) with wheat-based
 179 diets. A feed grain by CP interaction ($P = 0.006$) was observed for relative abdominal fat-pad weights.
 180 This was largely because birds offered 170 g/kg CP, sorghum-based diets had 40.1% heavier relative
 181 abdominal fat-pad weights (11.91 versus 8.50 g/kg) than their wheat-based counterparts.

182 As shown in Table 6, the dietary CP reduction decreased *Pectoralis major* yields by 6.22%
 183 (181 versus 193 g/kg) but did not significantly affect *Pectoralis minor* yields. The dietary CP reduction

184 increased leg quarter yields by 2.61% (236 versus 230 g/kg; $P = 0.021$) and expanded arginine:lysine
185 ratios slightly, but significantly decreased leg quarter yields.

186 Dietary treatment effects on protein and starch apparent ileal digestibility coefficients, disappearance
187 rates and starch to protein disappearance rate ratios are shown in Table 7. Sorghum-based diets
188 supported fractionally higher starch digestibility coefficients than wheat (0.996 versus 0.993; $P =$
189 0.034) and dietary CP reductions improved protein digestibility by 1.32% (0.845 versus 0.834; $P =$
190 0.042). A three-way treatment interaction ($P < 0.001$) was observed for starch disappearance rates
191 because expanded arginine:lysine ratios in 210 g/kg CP, sorghum-based diets significantly increased
192 disappearance rates by 11.4% (60.68 versus 54.49 g/bird per day). Dietary CP reductions depressed
193 protein disappearance rates by 18.8% (21.20 versus 26.10 g/bird per day; $P < 0.001$) and expanded
194 arginine:lysine ratios increased protein disappearance rates by 3.28% (23.92 versus 23.16 g/bird per
195 day; $P = 0.034$). A three-way treatment interaction ($P < 0.001$) was observed for starch to protein
196 disappearance rate ratios. Expanded arginine:lysine ratios in 210 g/kg CP, sorghum-based diets
197 significantly increased disappearance rate ratios from 2.12 to 2.28 and increased disappearance rate
198 ratios from 3.29 to 3.45 in 170 g/kg CP, wheat-based diets.

199 Dietary treatment effects on apparent ileal digestibility coefficients of essential amino acids
200 are shown in Table 8. A feed grain by CP interaction ($P = 0.001$) was observed for leucine because in
201 170 g/kg CP, wheat-based diets supported superior leucine digestibility by 7.63% (0.889 versus 0.826)
202 in comparison to sorghum-based diets. As a main effect, wheat generated significantly higher
203 digestibility coefficients for histidine (6.83%), isoleucine (2.92%), phenylalanine (3.06%) and valine
204 (2.30%) than sorghum, where the percentage increases are shown in parentheses. Alternatively,
205 sorghum supported higher digestibility coefficients for arginine (1.33%) and lysine (1.81%). Reducing
206 dietary CP levels significantly increased digestibilities of arginine (2.22%), isoleucine (5.54%), lysine
207 (3.08%), methionine (0.95%), phenylalanine (2.71%), threonine (4.05%) and valine (4.78%).

208 Increasing the dietary arginine:lysine ratio significantly depressed histidine and threonine
209 digestibilities by 1.58% and 1.95%, respectively.

210 The digestibility outcomes for non-essential amino acids are shown in Table 9. Feed grain by
211 CP interactions were observed for alanine, aspartic acid and proline ($P < 0.001$) and also tyrosine ($P =$
212 0.013). A weak three-way treatment interaction ($P = 0.024$) was observed for serine. The strong
213 interactions were driven by large differences in amino acid digestibilities between feed grains pursuant
214 to the dietary CP reduction. In 170 g/kg CP diets, sorghum supported noticeably higher mean
215 digestibilities for alanine (0.835 versus 0.678) and aspartic acid (0.783 versus 0.650); whereas, wheat
216 supported higher digestibilities for proline (0.911 versus 0.769) and tyrosine (0.871 versus 0.778). As
217 main effects, wheat-based diets generated higher glutamic acid digestibility by 8.50% (0.919 versus
218 0.847; $P < 0.001$) and the dietary CP reduction increased glycine digestibility by 9.92% (0.864 versus
219 0.786; $P < 0.001$).

220 The effects of dietary treatments on free plasma concentrations of essential and non-essential
221 amino acids are shown in Table 10 and Table 11, respectively. Feed grain by CP interactions were
222 observed for isoleucine ($P = 0.034$) and alanine ($P = 0.048$). With isoleucine, concentrations decreased
223 from 17.8 to 15.0 $\mu\text{g/g}$ following the dietary CP reductions in wheat-based diets, but increased from
224 11.6 to 14.0 $\mu\text{g/g}$ in sorghum-based diets. Concentrations of alanine decreased markedly from 106.8
225 to 67.7 $\mu\text{g/g}$ following the dietary CP reductions in wheat-based diets, but in sorghum-based diets the
226 decrease was relatively modest, from 106.5 to 95.1 $\mu\text{g/g}$. Wheat-based diets generated significantly
227 higher free plasma concentrations of histidine, valine, cysteine, glutamic acid, glutamine, glycine,
228 proline and serine; whereas, sorghum-based diets generated higher concentrations of arginine, leucine
229 and asparagine. Dietary CP reductions significantly increased concentrations of arginine (35.2%),
230 lysine (60.2%), methionine (78.4%), threonine (79.2%), valine (15.0%) and glycine (59.0%), but
231 decreased aspartic acid (24.5%) and cysteine (12.0%). Expanded arginine:lysine ratios significantly

232 increased arginine concentrations by 31.6% but decreased tryptophan by 17.7%. Concentrations of
233 phenylalanine and tyrosine were not statistically influenced by treatment.

234 4. Discussion

235 Overall growth performance in the present study was highly satisfactory as Ross 308 performance
236 objectives (Aviagen, 2019) for weight gain were exceeded by 18.6% (2193 versus 1849 g/bird) and
237 for FCR by 9.99% (1.423 versus 1.581). This is despite that reducing dietary CP by 40 g/kg
238 compromised weight gain by 10.9% (2078 versus 2310 g/bird) and FCR by 7.51% (1.474 versus 1.371)
239 as main effects. However, the dietary CP reduction depressed weight gain by 7.87% (2154 versus 2338
240 g/bird) and FCR by 4.03% (1.421 versus 1.366) in birds offered sorghum-based diets when average
241 dietary arginine:lysine ratios are combined. In contrast, growth performance of birds offered wheat-
242 based diets was compromised to greater extents with marked depressions of 12.2% (2002 versus 2280
243 g/bird) in weight gain and 11.5% (1.527 versus 1.378) in FCR. These data both reflect the challenges
244 to the successful development of reduced-CP diets and indicate that sorghum is a more suitable feed
245 grain than wheat in this context.

246 Instructively, NBAA inclusions were comparable (wheat: 9.80 g/kg, sorghum: 9.50 g/kg) in
247 210 g/kg CP diets, but in 170 g/kg CP diets NBAA inclusions were considerably higher in wheat-based
248 diets (48.96 versus 35.33 g/kg). The higher NBAA inclusions in wheat-based diets are driven by the
249 higher protein content of wheat (139 g/kg) than sorghum (107 g/kg) used in the present study, which
250 is a typical difference. However, Baker (2009) contended that there are limits to the extent that intact
251 protein can be replaced by non-bound amino acids to achieve maximal weight gain and feed efficiency.
252 If so, wheat is disadvantaged relative to sorghum in the framework of reduced-CP broiler diets.
253 Moreover, it may be deduced that increasing dietary NBAA inclusions were linearly associated with
254 less efficient weight gain ($r = -0.834$; $P < 0.001$) and FCR ($r = 0.862$; $P < 0.001$) in the present study.

255 While not conclusive, these relationships are consistent with the proposal that NBAA inclusions can
256 become excessive in reduced-CP diets.

257 Intestinal uptakes of NBAA are more rapid than their protein-bound counterparts (Wu, 2009)
258 and the implication is that non-bound and protein-bound amino acids are not bioequivalent (Selle et
259 al., 2022b). The likelihood is that this difference promotes post-enteral imbalances between non-bound
260 and protein-bound amino acids leading to post-prandial oxidation of surplus amino acids (Selle et al.,
261 2022b). For example, non-bound leucine was more susceptible to post-prandial oxidation than protein-
262 bound leucine in rats (Nolles et al., 2009). The catabolism of surplus amino acids is an obvious loss,
263 but it is accompanied by an 'energy cost' because an energy input of 60.7 kJ is required to eliminate 1
264 g of uric acid-N generated by amino acid catabolism (Van Milgen, 2021).

265 A three-way interaction ($P = 0.008$) between CP, feed grain and arginine:lysine ratio was
266 observed for FCR in the present study. In birds offered 170 g/kg CP sorghum-based, expanding
267 arginine:lysine ratios improved FCR by 2.30%, but depressed FCR by 2.12% in their wheat-based
268 counterparts. Arginine and lysine requirements for broiler chickens were determined by Nogueira et
269 al. (2022) and in male birds, optimal ratios ranged from 107 to 118 depending on age. However,
270 increasing dietary arginine:lysine ratios from 88 to 113 in maize-based diets improved FCR by 4.91%
271 (1.55 versus 1.63) in Castro et al. (2020). This parallels the response in sorghum-based diets in the
272 present study, but not wheat-based diets. In something of a precedent, elevated BCAA inclusions in
273 187.5 g/kg CP, wheat-based diets significantly depressed FCR by 8.33% (1.665 versus 1.537), but
274 fractionally improved FCR in sorghum-based diets (1.378 versus 1.390) in Greenhalgh et al. (2022).
275 Moreover, elevated BCAA inclusions decreased weight gain by 9.49% (1288 versus 1423 g/bird) in
276 wheat-based diets, but increased gain by 9.26% (1451 versus 1328 g/bird) in sorghum-based diets in
277 this study. Concentrations of non-bound BCAA, especially leucine, were substantially higher in
278 wheat- than sorghum-based diets in Greenhalgh et al. (2022) and these imbalances may have

279 contributed to the observed responses. In the present study, wheat-based diets contained more non-
280 bound arginine (16.50 versus 13.68 g/kg) and lysine (23.28 versus 19.29 g/kg) than sorghum and,
281 reciprocally, sorghum-based diets contained more protein-bound amino acids. These differences may
282 have exacerbated the recognised antagonism between arginine and lysine (Balnave and Brake, 2002);
283 the likely basis of this antagonism is that a relative excess of lysine may impede the renal re-
284 absorption of arginine (Maynard and Kidd, 2022). This may have contributed to the treatment
285 interaction observed for FCR in the present study.

286 The importance of considering starch and protein digestive dynamics in tandem was evident
287 in the present study as condensing ileal starch to protein disappearance rate ratios were quadratically
288 associated with improvements in weight gain ($r = 0.805$; $P < 0.001$) and FCR ($r = 0.780$; $P < 0.001$).
289 The positive impacts of capping dietary starch:protein ratios and, in turn, condensing starch to protein
290 disappearance rate ratios on growth performance of birds offered reduced-CP, wheat- and maize-based
291 diets has been previously reported (Greenhalgh et al., 2020; 2022b). Dietary starch:protein ratios will
292 typically expand in the formulation of reduced-CP diets and any strategies that will limit this trend
293 should be advantageous.

294 Perturbations in apparent amino acid digestibilities pursuant to dietary CP reductions are
295 commonly observed (Liu et al., 2021) and constitute an impediment to the precise formulation of
296 reduced-CP diets to meet amino acid requirements. The genesis of these perturbations is the opposing
297 forces that are in play. Average digestibilities of five amino acids (Ala, Asp, Glu, Pro, Ser) that were
298 present only as protein-bound entities in the present experiment decreased by 4.56% (0.690 versus
299 0.723) following the reduction from 210 to 170 g/kg CP. In contrast, average digestibilities of seven
300 amino acids (arginine, isoleucine, lysine, methionine, threonine, valine, glycine) that were included as
301 non-bound entities across all diets increased by 4.12% (0.885 versus 0.850). Dietary CP reductions can
302 reduce apparent amino acid digestibility coefficients because concentrations of dietary amino acids in

303 distal ileal digesta are diluted by amino acids derived from endogenous secretions and the gut
304 microbiota. This shift in amino acid concentrations depresses apparent digestibility coefficients
305 (Donkoh and Moughan, 1994). In addition, there are variations in inherent amino acid digestibilities
306 of the three key feedstuffs: soybean meal, wheat and sorghum. Ravindran et al. (1999) reported that
307 the mean ileal digestibility of 14 amino acids in soybean meal was 0.816 in comparison to 0.774 for
308 wheat and 0.743 for sorghum. Therefore, the partial substitution of soybean meal with either feed grain
309 in the formulation of reduced-CP diets will tend to depress amino acid digestibilities. Interestingly,
310 histidine digestibility in wheat was superior to sorghum by 12.4% (0.782 versus 0.696) in Ravindran
311 et al. (1999), which was reflected in the present study as wheat generated higher histidine digestibilities
312 than sorghum by 6.83% (0.845 versus 0.791) as a main effect. Theoretically, NBAA are completely
313 digestible (Lemme et al., 2005), which will counteract the above two negative factors when amino
314 acids are included in diet formulations as non-bound entities at high inclusion levels. For example, the
315 reduction in dietary CP increased lysine digestibility by 3.08% (0.903 versus 0.876; $P < 0.001$) where
316 lysine-HCl inclusions ranged from 3.55 to 10.70 g/kg in the present study.

317 Free amino acid concentrations in systemic plasma are difficult to interpret because they
318 reflect the dynamic equilibrium between post-enteral amino acid availability and protein accretion,
319 which is complicated by protein degradation, catabolism and gluconeogenesis involving amino acids
320 (Fernández-Fígares et al., 1997). Dietary CP reductions significantly increased free plasma
321 concentrations of methionine, lysine, threonine, valine, arginine; these increases could be indicative of
322 inefficient utilisation of these pivotal amino acids. Also, plasma concentrations of methionine ($r =$
323 0.625 ; $P < 0.001$), glycine ($r = 0.674$; $P < 0.001$), lysine ($r = 0.584$; $P < 0.001$), and threonine ($r =$
324 0.569 ; $P < 0.001$) were linearly related to increases in FCR or compromised feed efficiency. Again,
325 these positive relationships could be indicative of inefficient amino acid utilisation for protein
326 deposition. However, elevated plasma free threonine concentrations are frequently observed in broiler

327 chickens following dietary CP reductions and could even serve as a biomarker for the adequacy with
328 which reduced-CP diets are formulated (Macelline et al., 2021). Instructively, there is a negative linear
329 relationship ($r = -0.446$; $P = 0.004$) between the ratio of free arginine to lysine plasma concentrations
330 and FCR, as shown in Fig. 1. Thus, increases in arginine relative to lysine in the systemic circulation
331 was associated with enhanced FCR and the linear equation predicts that an increase in plasma ratios
332 from 1.0 to 2.0 would enhance FCR by 4.84% (1.377 versus 1.447). This may reflect post-enteral
333 antagonistic interactions between arginine and lysine antagonism (Kadirvel and Kratzer, 1974).

334 5. Conclusion

335 It was established that sorghum is a more suitable feed grain than wheat in reduced-CP broiler diets as
336 there was a CP \times feed grain interaction ($P = 0.013$); sorghum supported a 2.41% greater increase in
337 weight gain than wheat in 210 g/kg CP diets, but this advantage expanded to 7.59% in 170 g/kg CP
338 diets. Increasing dietary arginine:lysine ratios per se did not influence growth performance, but a three-
339 way FCR interaction ($P = 0.008$) showed that increasing arginine:lysine ratios in 170 g/kg CP,
340 sorghum-based diets generated a 2.30% improvement in FCR as opposed to a 2.12% depression in
341 FCR in corresponding wheat-based diets.

342 Acknowledgements

343 We would like to thank AgriFutures Chicken-meat for funding the project “Optimising amino acid
344 profiles and energy in reduced-protein diets” (PRJ-010520) and for their ongoing encouragement and
345 support. We would like to acknowledge the financial support provided by Australian Government
346 Research Training Program Scholarship (RTP) received by PhD candidate Mr Shemil Macelline. We
347 also acknowledge Ms. Joy Gill, Mr. Duwei Chen, Ms. Kylie Warr and Mr. Peter Bird from Poultry
348 Research Foundation for their assistance with feed manufacturing, bird management, sample
349 collection, and laboratory analyses.

350 References

- 351 Austic RE, Scott RL. Involvement of food intake in the lysine-arginine antagonism in chicks. *J Nutr*
352 1975;105:1122-31.
- 353 Aviagen. Ross 308/Ross 308 FF Broiler: Performance Objectives, 4. 2019. (Accessed on 09 June
354 2022) [http://eu.aviagen.com/tech-center/download/1339/Ross308-308FF-BroilerPO2019-](http://eu.aviagen.com/tech-center/download/1339/Ross308-308FF-BroilerPO2019-EN.pdf)
355 [EN.pdf](http://eu.aviagen.com/tech-center/download/1339/Ross308-308FF-BroilerPO2019-EN.pdf)
- 356 Bach Knudsen KE. Carbohydrate and lignin contents of plant materials used in animal feeding. *Anim*
357 *Feed Sci Tech* 1997;67:319–38.
- 358 Baker DH. Advances in protein-amino acid nutrition in poultry. *Amino Acids* 2009;37:29-41.
- 359 Ball RO, Urschel KL, Pencharz PB. Nutritional consequences of interspecies differences in arginine
360 and lysine metabolism. *J Nutr* 2007;137:1626–41.
- 361 Balnave D, Brake J. Re-evaluation of the classical dietary arginine:lysine interaction for modern
362 poultry diets: a review. *World's Poult Sci J* 2002;58:275–89.
- 363 Castro FLS, Kim WK. Secondary functions of arginine and sulfur amino acids in poultry health:
364 Review. *Animals* 2020;10:2106.
- 365 Castro LS, Teng P-Y, Yadav S, Gould RL, Craig S, Pazdro R, Kim WK. The effects of L-Arginine
366 supplementation on growth performance and intestinal health of broiler chickens challenged
367 with *Eimeria* spp. *Poult Sci* 2020;99:5844-57.
- 368 Chrystal PV, Greenhalgh S, McInerney BV, McQuade LR, Akter Y, de Paula Dorigam JC, Selle PH,
369 Liu SY. Maize-based diets are more conducive to crude protein reductions than wheat-based
370 diets for broiler chickens. *Anim Feed Sci Technol* 2021;275:114867.
- 371 Cohen SA, Michaud DP. Synthesis of a fluorescent derivatizing reagent, 6-aminoquinolyl-n-
372 hydroxysuccinimidyl carbamate, and its application for the analysis of hydrolysate amino-
373 acids via high-performance liquid-chromatography. *Anal Biochem* 1993;211:279–287.

- 374 Corzo A, Lee J, Vargas JI, Silva M, Pacheco WJ. Determination of the optimal digestible arginine to
375 lysine ratio in Ross 708 male broilers. *J Appl Poult Res* 2021;30:100136.
- 376 Donkoh A, Moughan PJ. The effect of dietary crude protein content on apparent and true ileal
377 nitrogen and amino acid digestibilities. *Brit Poult Sci* 1994;72:59-68.
- 378 Fernández-Fígares I, Prieto C, Nieto R, Aguilera JF (1997) Free amino acid concentrations in plasma,
379 muscle and liver as indirect measures of protein adequacy in growing chickens. *Anim Sci*
380 1997;64:1997;529-539.
- 381 Fouad AM, El-Senousey HK, Yang XJ, Yao JH. Dietary L -arginine supplementation reduces
382 abdominal fat content by modulating lipid metabolism in broiler chickens. *Animal*
383 2013;7:1239-45.
- 384 Giuberti G, Gallo A, Cerioli C, Masoero F. In vitro starch digestion and predicted glycemic index of
385 cereal grains commonly utilized in pig nutrition. *Anim Feed Sci Technol* 2012;174:163–73.
- 386 Gopinger E, Krabbe EL, Surek D, Lopes LS, Avila VS. Live performance, carcass, and bone quality
387 responses of grower and finisher broilers to dietary metabolizable energy levels. *Brazilian*
388 *Journal of Poult Sci.* 2017;19:559-66.
- 389 Greenhalgh S, McInerney BV, McQuade LR, Chrystal PV, Khoddami A, Zhuang MAM, Liu SY,
390 Selle PH. Capping dietary starch:protein ratios in moderately reduced crude protein, wheat-
391 based diets showed promise but further reductions generated inferior growth performance in
392 broiler chickens. *Anim Nutr* 2020;6:168-78.
- 393 Greenhalgh S, Macelline SP, Chrystal PV, Liu SY, Selle PH. An evaluation of elevated branched-
394 chain amino acid inclusions on the performance of broiler chickens offered reduced-crude
395 protein, wheat-based diets from 7 to 28 days post-hatch. *Anim Feed Sci Technol*
396 2022a;286:115255.

- 397 Greenhalgh S, Chrystal PV, Lemme A, Dorigam JC, Macelline SP, Liu SY, Selle PH. Capping dietary
398 starch: Protein ratios enhances performance of broiler chickens offered reduced-crude protein,
399 maize-based diets. *Anim Feed Sci Technol* 2022b;290:115374.
- 400 Kadirvel R, Kratzer FH. Uptake of l-arginine and l-lysine by the small intestine and its influence on
401 arginine-lysine antagonism in chicks. *J Nutr* 1974;104:339-343.
- 402 Lemme A, Rostagno HS, Petri A, Albino LF. Standardised ileal digestibility of crystalline amino
403 acids. *Proc 15th Euro Symp Poult Nutr* 2005;462–464. Balatonfüred, Hungary. WPSA.
404 <https://www.cabdirect.org/cabdirect/abstract/20073279556>
- 405 Liu SY, Macelline SP, Chrystal PV, Selle PH. Progress towards reduced-crude protein diets for
406 broiler chickens and sustainable chicken-meat production. *J Anim Sci Biotechnol* 2021;12:
407 20.
- 408 Macelline SP, Chrystal PV, Lu SY, Selle PH. Implications of elevated threonine plasma
409 concentrations in the development of reduced-crude protein diets for broiler chickens. *Anim*
410 *Prod Sci* 2021;61:1442-1448.
- 411 Mahasukhonthachat, K, Sopade P, Gidley M. Kinetics of starch digestion and functional properties
412 of twin-screw extruded sorghum. *J Cereal Sci* 2010;51:392-401.
- 413 Maynard CW, Kidd MT. Broiler Amino Acid Research: Then and Now. In: Téllez G, Latorre JD,
414 Martínez-Aguilar PY editors. *Broiler Industry*. London: IntechOpen; 2022. Available from:
415 <https://www.intechopen.com/online-first/79974> doi: 10.5772/intechopen.101896
- 416 Nogueira RF, Sakomura NKL, Leme BB, de Paula Reis M, Kochenborger JB, Fernandes JBK, da
417 Silva Viana G (2022) Lysine and arginine requirements of male and female broiler in the
418 starter, grower, and finisher phase. *Anim Feed Sci Technol* 2022;283:115174.

- 419 Nolles JA, Verreijen AM, Koopmanschap RE, Verstegen MWA, Schreurs VVAM. Postprandial
420 oxidative losses of free and protein-bound amino acids in the diet: interactions and adaptation.
421 *J Anim Physiol Anim Nutr* 2009;93:431-38.
- 422 OECD Data. Meat consumption (indicator).doi:10.1787/fa290fd0-en.
423 <https://data.oecd.org/agrooutput/meat-consumption.htm> [Accessed on 09 June 2022]
- 424 Ravindran V, Cabahug S, Ravindran G, Bryden WL. Influence of microbial phytase on apparent ileal
425 amino acid digestibility of feedstuffs for broilers. *Poult Sci* 1999;78:699-706.
- 426 Selle PH, Macelline SP, Greenhalgh S, Chrystal PV, Liu SY. Identifying the shortfalls of crude
427 protein-reduced, wheat-based broiler diets. *Anim Nutr* 2022a;11:181-189.
- 428 Selle PH, Macelline SP, Chrystal PV, Liu SY. The impact of digestive dynamics on the
429 bioequivalence of amino acids in broiler chickens. *Frontiers in Bioscience – Landmark*
430 2022b;27:126.
- 431 Siriwan P, Bryden WL, Mollah Y, Annison EF. Measurement of endogenous amino acid losses in
432 poultry. *Brit Poult Sci* 1993;34:939-49.
- 433 Van Milgen J. The role of energy, serine, glycine, and 1-carbon units in the cost of nitrogen excretion
434 in mammals and birds. *Animal* 2021;5:100213.
- 435 Wu G. Amino acids: metabolism, functions, and nutrition. *Amino Acids* 2014;37:1-17.
- 436 Wu G. Dietary requirements of synthesizable amino acids by animals: a paradigm shift in protein
437 nutrition. *J Anim Sci Biotech* 2014;5:34.
- 438 Yin D, Chrystal PV, Moss AF, Liu SY, Yian J, Selle PH. Effects of reducing dietary crude protein
439 and whole grain feeding on performance and amino acid metabolism in broiler chickens
440 offered wheat-based diets. *Anim Feed Sci Technol* 2020;260:114386.

441 Zampiga M, Laghi L, Petracci M, Zhu C, Meluzzi A, Dridi S, Sirri F Effect of dietary arginine to
442 lysine ratios on productive performance, meat quality, plasma and muscle metabolomics
443 profile in fast-growing broiler chickens. J Anim Sci Biotech 2018;9:79.

Journal Pre-proof

Table 1. Outline of 8 dietary treatments

Experimental diet	Crude protein, g/kg	Feed grain	Arginine:lysine ratio
1A	210	Wheat	104
2B	210	Sorghum	104
3C	170	Wheat	104
4D	170	Sorghum	104
5E	210	Wheat	110
6F	210	Sorghum	110
7G	170	Wheat	110
8H	170	Sorghum	110

Table 2. Composition of experimental diets (g/kg, as-is basis)

Item	1A	2B	3C	4D	5E	6F	7G	8H
Sorghum	-	624	-	789	-	623	-	794
Wheat	674	-	877	-	679	-	863	-
Soybean meal	225	275	-	102	219	275	-	95.8
D,L-Methionine	2.68	3.21	4.31	4.66	2.72	3.21	4.38	4.72
Glycine	0.13	0.98	5.57	5.63	0.28	0.98	5.7	5.8
L-Arginine	0.69	0.56	6.77	5.52	1.53	1.23	7.51	6.37
L-Histidine	-	-	1.25	1.04	-	-	1.29	1.1
L-Isoleucine	0.44	0.13	3.82	2.82	0.54	0.14	3.89	2.92
L-Leucine	-	-	4.84	-	-	-	4.96	-
L-Lysine HCl	4.2	3.55	10.7	8.72	4.38	3.55	10.7	8.91
L-Phenylalanine	-	-	2.53	1.26	-	-	2.53	1.36
L-Threonine	1.68	1.35	4.53	3.6	1.76	1.35	4.58	3.68
L-Tryptophan	-	-	0.65	0.24	-	-	0.67	0.27
L-Tyrosine	-	-	2.14	0.59	-	-	2.26	0.69
L-Valine	0.9	0.5	4.3	3.17	1	0.5	4.38	3.27
Soy oil	43.9	45.1	12.3	19.6	43.1	45.1	17	18.7
Limestone	13.8	13.6	14.8	14.4	13.9	13.6	14.8	14.4
Monocalcium phosphorus	6.5	6.16	8.08	7.34	6.54	6.16	8.14	7.39
Potassium carbonate	-	-	6.44	2.62	-	-	6.51	2.85
Salt	1.38	3	-	-	1.19	3	-	-
Sodium bicarbonate	2.24	-	4.33	4.45	2.52	-	4.33	4.46
Vitamin-mineral premix ¹	2	2	2	2	2	2	2	2
Choline Cl (60%)	0.8	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Feed enzymes ²	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Celite	20	20	20	20	20	20	20	20
Inert filler	-	-	2.37	-	-	-	10.8	-

¹The vitamin-mineral premix supplied per tonne of feed: retinol, 12,000 IU; cholecalciferol, 5000 IU; tocopheryl acetate, 75 mg; menadione, 3 mg; thiamine, 3 mg; riboflavin, 8 mg; niacin, 55 mg; pantothenate, 13 mg; pyridoxine, 5 mg; folate, 2 mg; cyanocobalamin, 16 µg; biotin, 200 µg; cereal-based carrier, 149 mg; Cu (sulphate), 16 mg; Fe (sulphate), 40 mg; I (iodide), 1.25 mg; Se (selenate), 0.3 mg; Mn (sulphate and oxide), 120 mg; Zn (sulphate and oxide), 100 mg; cereal-based carrier, 128 mg;

²Phytase (Aextra PHY, Danisco Animal Nutrition) and xylanase (Aextra XB, Danisco Animal Nutrition)

Table 3. Nutrient specifications of experimental diets (g/kg)

Item	1A	2B	3C	4D	5E	6F	7G	8H
AME, MJ/kg	13.0	13.0	13.0	13.0	13.0	13.0	13.0	13.0
Crude protein	210	210	170	170	210	210	170	170
SID lysine	11.0	11.0	11.0	11.0	11.0	11.0	11.0	11.0
SID methionine	5.21	5.77	5.89	6.45	5.23	5.77	5.92	6.48
SID TSAA	8.14	8.14	8.14	8.14	8.14	8.14	8.14	8.14
SID threonine	7.26	7.26	7.26	7.26	7.26	7.26	7.26	7.26
SID valine	8.69	8.69	8.69	8.69	8.69	8.69	8.69	8.69
SID isoleucine	7.59	7.59	7.59	7.59	7.59	7.59	7.59	7.59
SID leucine	12.4	16.4	11.8	13.1	13	16.4	11.8	12.9
SID arginine	11.4	11.4	11.4	11.4	12.1	12.1	12.1	12.1
SID histidine	4.28	4.18	3.63	3.63	4.23	4.17	3.63	3.63
SID tryptophan	2.15	2.31	1.76	1.76	2.12	2.31	1.76	1.76
SID glycine equivalent ¹	13.1	13.1	13.1	13.1	13.1	13.1	13.1	13.1
SID phenylalanine	8.54	8.96	4.84	6.06	8.43	8.96	4.76	5.95
SID Phenylalanine + tyrosine	13.5	14.8	11.6	11.6	13.4	14.8	11.6	11.6
Calcium	8.70	8.70	8.70	8.70	8.70	8.70	8.70	8.70
Available phosphorus	4.35	4.35	4.35	4.35	4.35	4.35	4.35	4.35
Crude fibre	20.4	20.4	17.6	18.2	20.4	20.4	17.3	18.1
Crude fat	61.5	72.5	30.5	50.4	60.7	72.5	34.7	49.5
DEB	210	210	210	210	210	210	210	210

SID = standard ileal digestibility; TSAA = total sulphur amino acids; DEB = dietary electrolyte balance = $K^+ + Na^+ - Cl^-$

¹ Glycine equivalent = glycine concentration + (serine concentration \times 0.7143).

Table 4. Analysed nutrient composition of experimental diets (as fed-basis, g/kg)

Item	1A	2B	3C	4D	5E	6F	7G	8H
Dry matter	893	880	890	875	893	883	894	871
Crude protein	205	204	170	168	210	210	171	180
Starch	374	365	483	497	380	400	500	478
Arginine	11.5	11.7	10.9	10.9	12.7	11.5	11.7	12.0
Histidine	5.10	5.30	4.00	4.20	5.20	5.00	3.80	4.20
Isoleucine	8.40	9.20	7.40	8.30	8.90	8.90	7.40	8.60
Leucine	14.0	19.4	12.3	15.7	14.6	19.5	12.3	15.8
Lysine	11.6	12.1	11.1	10.9	12.5	11.3	11.1	11.6
Methionine	4.30	4.30	5.00	5.10	4.10	4.00	5.30	5.50
Phenylalanine	9.40	10.3	6.90	8.30	9.80	10.1	7.80	8.50
Threonine	8.10	8.50	7.40	7.60	8.50	8.10	7.40	7.90
Valine	9.80	10.3	9.20	9.50	10.3	10.0	9.30	9.80
Alanine	7.40	8.50	4.00	9.90	7.80	11.6	3.90	9.80
Aspartic acid	16.4	19.5	6.00	11.1	17.4	18.1	5.80	11.0
Glutamic acid	46.7	39.0	36.7	28.0	49.1	38.0	35.4	27.9
Glycine	8.20	8.50	10.2	9.50	8.70	8.10	10.7	10.1
Proline	14.3	12.1	13.0	9.50	14.8	12.2	11.8	9.70
Serine	9.30	9.60	5.50	6.30	9.70	9.40	5.40	6.40
Tyrosine	4.10	4.30	3.60	3.10	4.10	3.90	3.70	3.10
Total amino acids	189	196	153	158	198	190	153	162

Table 5. Effects of dietary treatments on growth performance and relative abdominal fat pad weights from 14 to 35 d post-hatch

Crude protein, g/kg	Treatment		Growth performance			Relative fat-pad weights, g/kg
	Feed grain	Arginine:lysine ratio	Weight gain, g/kg	Feed intake, g/bird	FCR, g/g	
210	Wheat	104	2252 ^c	3123	1.387 ^{ab}	8.09 ^a
		110	2308 ^{cd}	3161	1.369 ^a	8.65 ^{ab}
	Sorghum	104	2332 ^d	3192	1.369 ^a	9.65 ^{bc}
		110	2343 ^d	3192	1.362 ^a	10.15 ^c
170	Wheat	104	2024 ^a	3058	1.511 ^d	8.55 ^{ab}
		110	1979 ^a	3053	1.543 ^c	8.45 ^{ab}
	Sorghum	104	2147 ^b	3085	1.437 ^c	12.15 ^d
		110	2161 ^b	3034	1.404 ^b	11.67 ^d
SEM			26.0	36.7	0.0097	0.043
Main effects: FG						
	Wheat		2137	3098	1.455	8.45
	Sorghum		2243	3128	1.394	10.95
CP, g/kg						
210			2310	3168 ^b	1.371	9.15
170			2078	3057 ^a	1.474	10.21
Arginine:lysine ratio						
	104		2181	3111	1.423	9.66
	110		2198	3110	1.420	9.73
Significance (<i>P</i> -value)						
	FG		<0.001	0.302	<0.001	< 0.001
	CP		<0.001	<0.001	<0.001	0.002
	Arginine:lysine ratio		0.623	0.868	0.345	0.710
	FG × CP		0.013	0.382	<0.001	0.006
	FG × arginine:lysine ratio		0.840	0.423	0.058	0.742
	CP × arginine:lysine ratio		0.179	0.375	0.375	0.219
	FG × CP × Arginine:lysine ratio		0.168	0.936	0.008	0.807

FG = feed grain; CP = crude protein.

^{a b c d} Within a column, means without a common superscript differ at $P < 0.05$.

Table 6. Effects of dietary treatments on relative weights (g/kg) of carcass traits at day 35 post-hatch

Crude protein, g/kg	Treatment		<i>Pectoralis major</i>	<i>Pectoralis minor</i>	Leg quarters
	Feed grain	Arginine:lysine ratio			
210	Wheat	104	204	34.7	223
		110	181	31.7	233
	Sorghum	104	193	31.8	229
		110	194	28.7	234
170	Wheat	104	187	31.3	232
		110	186	32.1	238
	Sorghum	104	179	30.5	234
		110	172	30.8	241
SEM			5.9	2.11	3.8
Main effects: FG					
	Wheat		189	32.5	231
	Sorghum		184	30.5	234
CP, g/kg					
210			193 ^b	31.7	230 ^a
170			181 ^a	31.2	236 ^b
Arginine:lysine ratio					
		104	191	32.1	229 ^b
		110	183	30.8	227 ^a
Significance (<i>P</i> -value)					
			0.247	0.191	0.282
			0.007	0.715	0.021
			0.087	0.402	0.012
			0.169	0.524	0.901
			0.295	0.928	0.698
			0.389	0.237	0.844
			0.105	0.954	0.454

FG = feed grain; CP = crude protein.

^{a,b} Within a column, means without a common superscript differ at $P < 0.05$.

Table 7. Effects of dietary treatments on protein and starch digestibility coefficients, disappearance rates and starch to protein disappearance rate ratios in distal ileum at 35 d post-hatch

Crude protein, g/kg	Treatment		Digestibility coefficients		Disappearance rates, g/bird per day		Starch to protein disappearance rate ratios	
	Feed grain	Arginine:lysine ratio	Starch	Protein	Starch	Protein		
210	Wheat	104	0.995	0.844	55.34 ^a	25.72	2.15 ^a	
		110	0.995	0.837	56.92 ^a	26.45	2.16 ^a	
	Sorghum	104	0.994	0.830	54.49 ^a	25.75	2.12 ^a	
		110	0.997	0.826	60.68 ^b	26.35	2.28 ^b	
170	Wheat	104	0.992	0.859	69.79 ^{cd}	21.26	3.29 ^d	
		110	0.989	0.840	71.98 ^{de}	20.89	3.45 ^e	
	Sorghum	104	0.997	0.836	72.74 ^c	20.64	3.53 ^e	
		110	0.995	0.846	69.01 ^c	21.99	3.13 ^c	
		SEM		0.0018	0.0075	0.880	0.367	0.029
		Main effects: FG						
Wheat			0.993 ^a	0.845	64.07	23.50	2.81	
Sorghum			0.996 ^b	0.835	63.96	23.61	2.75	
CP, g/kg								
210			0.995	0.834 ^a	58.86	26.10 ^b	2.18	
170			0.993	0.845 ^b	71.19	21.20 ^a	3.38	
Arginine:lysine ratio								
104			0.995	0.843	63.72	23.16 ^a	2.82	
110			0.994	0.837	64.38	23.92 ^b	2.74	
Significance (<i>P</i> -value)								
FG			0.034	0.060	0.259	0.702	0.884	
CP			0.150	0.042	<0.001	<0.001	<0.001	
Arginine:lysine ratio			0.693	0.326	0.018	0.034	0.523	
FG × CP			0.079	0.714	0.250	0.802	0.054	
FG × arginine:lysine ratio			0.580	0.159	0.607	0.137	<0.001	
CP × arginine:lysine ratio			0.183	0.918	0.001	0.731	<0.001	
FG × CP × arginine:lysine ratio			0.838	0.235	<0.001	0.088	<0.001	

FG = feed grain; CP = crude protein.

^{a b c d} Within a column, means without a common superscript differ at $P < 0.05$.

Table 8. Effects of dietary treatments on apparent digestibility coefficients of essential amino acids in distal ileum at 35 d post-hatch

Crude protein, g/kg	Treatment		Arginine	Histidine	Isoleucine	Leucine	Lysine	Methionine	Phenylalanine	Threonine	Valine
	Feed grain	Arginine:lysine ratio									
210	Wheat	104	0.896	0.842	0.843	0.845 ^b	0.874	0.946	0.865	0.801	0.829
		110	0.892	0.831	0.839	0.837 ^{ab}	0.869	0.949	0.859	0.794	0.822
	Sorghum	104	0.903	0.798	0.826	0.831 ^{ab}	0.888	0.953	0.844	0.792	0.814
		110	0.904	0.788	0.811	0.817 ^a	0.875	0.946	0.834	0.776	0.800
170	Wheat	104	0.918	0.865	0.897	0.897 ^c	0.899	0.957	0.900	0.851	0.876
		110	0.905	0.840	0.881	0.881 ^c	0.885	0.951	0.884	0.819	0.855
	Sorghum	104	0.925	0.792	0.865	0.830 ^{ab}	0.915	0.960	0.859	0.827	0.847
		110	0.927	0.789	0.861	0.822 ^{ab}	0.912	0.963	0.853	0.823	0.844
SEM			0.0049	0.062	0.0080	0.0092	0.0067	0.0037	0.0075	0.0087	0.0083
Main effects: FG											
	Wheat		0.903 ^a	0.845 ^b	0.866 ^b	0.866	0.882 ^a	0.952	0.875 ^b	0.817	0.846 ^b
	Sorghum		0.915 ^b	0.791 ^a	0.841 ^a	0.825	0.898 ^b	0.954	0.849 ^a	0.805	0.827 ^a
CP, g/kg											
	210		0.899 ^a	0.814	0.830 ^a	0.832	0.876 ^a	0.948 ^a	0.850 ^a	0.791 ^a	0.816 ^a
	170		0.919 ^b	0.822	0.876 ^b	0.858	0.903 ^b	0.957 ^b	0.873 ^a	0.823 ^b	0.855 ^b
Arginine:lysine ratio											
	104		0.911	0.825 ^b	0.860	0.852	0.895	0.954	0.864	0.819 ^b	0.843
	110		0.907	0.812 ^a	0.848	0.839	0.885	0.952	0.860	0.803 ^a	0.830
Significance (<i>P</i> -value)											
	FG		0.001	<0.001	<0.001	<0.001	0.002	0.348	<0.001	0.066	0.002
	CP		<0.001	0.224	<0.001	<0.001	<0.001	0.001	<0.001	<0.001	<0.001
	Arginine:lysine ratio		0.337	0.027	0.091	0.092	0.069	0.584	0.070	0.022	0.055
	FG × CP		0.518	0.097	0.759	0.001	0.233	0.078	0.218	0.717	0.874
	FG × arginine:lysine ratio		0.163	0.291	0.926	0.999	0.861	0.084	0.732	0.416	0.650
	CP × arginine:lysine ratio		0.562	0.785	0.977	0.945	0.953	0.883	0.789	0.595	0.842
	FG × CP × arginine:lysine ratio		0.539	0.333	0.335	0.585	0.260	0.875	0.478	0.142	0.281

FG = feed grain; CP = crude protein.

^{a b c} Within a column, means without a common superscript differ at $P < 0.05$.

Table 9. Effects of dietary treatments on apparent digestibility coefficients of non-essential amino acids in distal ileum at 35 d post-hatch

Crude protein, g/kg	Treatment		Alanine	Aspartic acid	Glutamic acid	Glycine	Proline	Serine	Tyrosine
	Feed grain	Arginine:lysine ratio							
210	Wheat	104	0.783 ^c	0.783 ^c	0.913	0.790	0.892 ^c	0.824 ^{de}	0.839 ^{de}
		110	0.784 ^c	0.787 ^c	0.913	0.783	0.891 ^c	0.826 ^e	0.828 ^{cd}
170	Sorghum	104	0.823 ^c	0.802 ^c	0.858	0.791	0.787 ^b	0.813 ^{de}	0.800 ^{bc}
		110	0.801 ^c	0.776 ^c	0.845	0.780	0.782 ^b	0.795 ^{bcd}	0.758 ^a
	Wheat	104	0.704 ^b	0.688 ^b	0.931	0.878	0.923 ^d	0.808 ^{cde}	0.878 ^f
		110	0.651 ^a	0.632 ^a	0.919	0.853	0.899 ^c	0.762 ^a	0.863 ^{ef}
Sorghum	104	0.825 ^c	0.784 ^c	0.846	0.862	0.753 ^a	0.778 ^{ab}	0.780 ^{ab}	
	110	0.819 ^c	0.782 ^c	0.841	0.864	0.754 ^a	0.782 ^{abc}	0.776 ^{ab}	
SEM			0.0158	0.0142	0.007	0.0082	0.0078	0.0106	0.0104
Main effects: FG									
	Wheat		0.729	0.720	0.919 ^b	0.827	0.902	0.804	0.852
	Sorghum		0.817	0.785	0.847 ^a	0.826	0.768	0.791	0.778
CP, g/kg									
	210		0.797	0.786	0.882	0.786 ^a	0.838	0.814	0.805
	170		0.750	0.721	0.884	0.864 ^b	0.833	0.782	0.824
Arginine:lysine ratio									
	104		0.783	0.762	0.887	0.833	0.839	0.805	0.825
	110		0.764	0.744	0.879	0.820	0.832	0.791	0.806
Significance (<i>P</i> -value)									
	FG		<0.001	<0.001	<0.001	0.803	<0.001	0.089	<0.001
	CP		<0.001	<0.001	0.699	<0.001	0.344	<0.001	0.020
	Arginine:lysine ratio		0.083	0.054	0.143	0.079	0.198	0.060	0.021
	FG × CP		<0.001	<0.001	0.051	0.893	<0.001	0.296	0.013
	FG × arginine:lysine ratio		0.605	0.562	0.799	0.318	0.364	0.329	0.505
	CP × arginine:lysine ratio		0.398	0.389	0.815	0.864	0.460	0.407	0.245
	FG × CP × arginine:lysine ratio		0.132	0.045	0.324	0.188	0.204	0.024	0.160

FG = feed grain; CP = crude protein.

^{a b c d} Within a column, means without a common superscript differ at $P < 0.05$.

Table 10. Effects of dietary treatments on essential amino acid plasma concentrations ($\mu\text{g/g}$) in broiler chickens

Crude protein, g/kg	Treatment		Arginine	Histidine	Isoleucine	Leucine	Lysine	Methionine	Phenylalanine	Threonine	Tryptophan	Valine
	Feed grain	Arginine:lysine ratio										
210	Wheat	104	39.5	8.1	19.3 ^b	28.0	37.0	23.6	24.8	92.5	6.5	41.1
		110	53.2	7.6	16.2 ^{ab}	23.6	35.2	21.4	20.3	72.1	5.3	35.6
	Sorghum	104	43.7	6.4	11.6 ^a	31.5	42.6	19.0	21.5	61.4	6.7	24.0
		110	61.9	7.0	11.6 ^a	32.6	33.9	19.0	23.3	66.7	5.4	24.5
170	Wheat	104	50.7	6.2	15.6 ^{ab}	24.4	67.9	38.5	23.6	139.8	6.9	41.2
		110	63.9	5.2	14.3 ^a	23.0	58.1	39.0	22.9	132.3	5.2	38.8
	Sorghum	104	68.7	5.4	14.2 ^a	29.2	58.1	35.2	25.2	127.1	5.0	32.6
		110	87.6	5.3	13.8 ^a	28.9	54.5	35.9	24.7	125.5	4.5	31.4
SEM			8.17	0.71	1.66	2.43	5.68	3.26	2.19	14.00	3.25	3.25
Main effects: FG												
	Wheat		51.8 ^a	6.7 ^b	16.4	24.8 ^a	49.6	30.6	22.9	109.2	5.9	39.2 ^b
	Sorghum		65.5 ^b	6.0 ^a	12.8	30.6 ^b	47.3	27.3	23.7	95.2	5.4	28.1 ^a
CP, g/kg												
	210		50.0 ^a	7.3	14.7	28.9	37.2 ^a	20.8 ^a	22.5	73.2 ^a	5.9	31.3 ^a
	170		67.6 ^b	5.5	14.5	26.4	59.6 ^b	37.1 ^b	24.1	131.2 ^b	5.4	36.0 ^b
Arginine:lysine ratio												
	104		50.6 ^a	6.5	15.2	28.3	51.4	29.1	23.8	105.2	6.2 ^b	34.7
	110		66.6 ^b	6.3	14.0	27.0	45.4	28.9	22.8	99.2	5.1 ^a	32.6
Significance (<i>P</i> -value)												
	FG		0.029	0.001	0.005	0.002	0.574	0.161	0.618	0.166	0.252	<0.001
	CP		0.004	0.158	0.871	0.149	<0.001	<0.001	0.301	<0.001	0.254	0.049
	Arginine:lysine ratio		0.010	0.614	0.317	0.473	0.147	0.935	0.535	0.547	0.023	0.352
	FG × CP		0.219	0.437	0.034	0.799	0.279	0.965	0.544	0.672	0.150	0.195
	FG × arginine:lysine ratio		0.657	0.516	0.401	0.349	0.967	0.788	0.287	0.429	0.567	0.427
	CP × arginine:lysine ratio		0.995	0.357	0.782	0.817	0.861	0.745	0.819	0.881	0.836	0.878
	FG × CP × arginine:lysine ratio		0.952	0.901	0.632	0.545	0.423	0.808	0.333	0.620	0.467	0.611

FG = feed grain; CP = crude protein.

^{a,b} Within a column, means without a common superscript differ at $P < 0.05$.

Table 11. Effects of dietary treatments on non-essential amino acid plasma concentrations ($\mu\text{g/g}$) in broiler chickens

Crude protein, g/kg	Treatment		Alanine	Asparagine	Aspartic acid	Cysteine	Glutamic acid	Glutamine	Glycine	Proline	Serine	Tyrosine
	Feed grain	Arginine:lysine ratio										
210	Wheat	104	122.0 ^c	37.5	17.3	19.6	31.9	411.0	72.2	87.8	81.1	59.0
		110	91.6 ^{bc}	30.9	13.1	16.5	27.6	337.0	55.7	77.9	63.6	44.0
	Sorghum	104	106.9 ^{bc}	33.5	15.8	13.4	24.0	257.0	58.0	50.6	62.2	55.5
		110	106.0 ^{bc}	37.2	10.9	13.8	22.8	232.0	57.1	53.7	62.3	55.1
170	Wheat	104	71.8 ^{ab}	18.7	9.3	15.3	23.8	375.0	98.4	72.9	73.0	51.6
		110	63.5 ^a	20.6	8.4	15.3	25.4	360.0	109.0	74.6	75.6	51.5
	Sorghum	104	98.2 ^{abc}	29.6	12.6	12.0	23.8	298.0	93.5	48.6	61.7	54.3
		110	92.0 ^{bc}	28.1	12.9	12.9	24.0	265.0	85.1	46.8	59.6	56.4
SEM			9.56	4.22	2.23	1.34	2.13	31.70	7.21	7.28	7.03	5.47
Main effects: FG												
	Wheat		87.2	26.9 ^a	12.0	16.6 ^b	27.2 ^b	371 ^b	83.6 ^b	78.3 ^b	73.3 ^b	51.5
	Sorghum		100.8	32.1 ^b	13.1	13.0 ^a	23.7 ^a	263 ^a	73.4 ^a	49.9 ^a	61.4 ^a	55.3
CP, g/kg												
210			106.3	34.8	14.3 ^b	15.8 ^b	26.6	309	60.7 ^a	67.5	67.3	53.4
170			81.4	24.3	10.8 ^a	13.9 ^a	24.3	324	96.5 ^b	60.7	67.4	53.5
Arginine:lysine ratio												
	104		99.7	29.8	13.8	15.1	25.9	335.0	80.5	65.0	69.5	55.1
	110		88.3	29.2	11.3	14.6	25.0	299.0	76.7	63.3	65.2	51.8
Significance (<i>P</i> -value)												
	FG		0.054	0.001	0.515	0.001	0.025	<0.001	0.050	<0.001	0.023	0.327
	CP		0.001	0.093	0.034	0.047	0.133	0.502	<0.001	0.194	0.968	0.987
	Arginine:lysine ratio		0.100	0.837	0.134	0.646	0.555	0.113	0.457	0.740	0.397	0.350
	FG × CP		0.048	0.190	0.081	0.400	0.071	0.345	0.437	0.656	0.722	0.993
	FG × arginine:lysine ratio		0.253	0.781	0.942	0.253	0.775	0.732	0.870	0.650	0.378	0.282
	CP × arginine:lysine ratio		0.539	0.559	0.183	0.347	0.240	0.565	0.342	0.746	0.523	0.270
	FG × CP × arginine:lysine ratio		0.321	0.258	0.785	0.491	0.446	0.454	0.100	0.428	0.272	0.424

FG = feed grain; CP = crude protein.

^{a b c} Within a column, means without a common superscript differ at $P < 0.05$.




Fig. 1 Negative linear relationship ($r = -0.446$; $P = 0.004$): $y = 1.517 - 0.070 x$, where y is FCR and x is the ratio of arginine to lysine plasma concentrations.

Conflict of interest

We declare that we have no financial and personal relationships with other people or organizations that can inappropriately influence our work, there is no professional or other personal interest of any nature or kind in any product, service and/or company that could be construed as influencing the content of this paper.

Journal Pre-proof