

Journal Pre-proof



No evidence of SARS-CoV-2 in semen of males recovering from COVID-19

Feng Pan, MD, Xingyuan Xiao, MD, Jingtao Guo, PhD, Yarong Song, MD, Honggang Li, MD, Darshan P. Patel, MD, Adam M. Spivak, MD, Joseph, P. Alukal, MD, Xiaoping Zhang, MD, Chengliang Xiong, MD, Philip S. Li, MD, James M. Hotaling, MD, MS

PII: S0015-0282(20)30384-8

DOI: <https://doi.org/10.1016/j.fertnstert.2020.04.024>

Reference: FNS 32440

To appear in: *Fertility and Sterility*

Received Date: 10 April 2020

Revised Date: 12 April 2020

Accepted Date: 13 April 2020

Please cite this article as: Pan F, Xiao X, Guo J, Song Y, Li H, Patel DP, Spivak AM, Alukal J,P, Zhang X, Xiong C, Li PS, Hotaling JM, No evidence of SARS-CoV-2 in semen of males recovering from COVID-19, *Fertility and Sterility* (2020), doi: <https://doi.org/10.1016/j.fertnstert.2020.04.024>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright ©2020 Published by Elsevier Inc. on behalf of the American Society for Reproductive Medicine

Running Title: No SARS-CoV-2 in semen of COVID-19 males

Title: No evidence of SARS-CoV-2 in semen of males recovering from COVID-19

Feng Pan, MD^{1,*}, Xingyuan Xiao, MD^{1,*}, Jingtao Guo, PhD^{2,3,*}, Yarong Song, MD^{1,*}, Honggang Li, MD^{4,*}, Darshan P. Patel, MD², Adam M. Spivak, MD⁵, Joseph, P. Alukal, MD⁶, Xiaoping Zhang, MD¹, Chengliang Xiong, MD^{7,8}, Philip S. Li, MD⁹, James M. Hotaling, MD, MS².

¹Department of Urology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China.

²Division of Urology, Department of Surgery, University of Utah School of Medicine, Salt Lake City, UT, USA.

³Department of Oncological Sciences, Huntsman Cancer Institute, Salt Lake City, UT, USA.

⁴Institute of Reproductive Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, China

⁵Division of Infectious Disease, Department of Internal Medicine, University of Utah School of Medicine, Salt Lake City, UT, USA.

⁶Department of Urology, Columbia University Medical Center, New York, NY, USA.

⁷Center for Reproductive Medicine, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430013, China

⁸Wuhan Tongji Reproductive Medicine Hospital, Wuhan, 43013, China.

⁹Center for Male Reproductive Medicine and Microsurgery, Department of Urology, Weill Cornell Medicine of Cornell University, New York, NY, USA.

*Similar in author order

Email Addresses for all authors:

Feng Pan, MD, panfeng@hust.edu.cn
Xingyuan Xiao, MD, xiaoxy@hust.edu.cn
Jingtao Guo, PhD, Jingtao.Guo@hci.utah.edu
Yarong Song, MD, song_yr@hust.edu.cn
Darshan P. Patel, MD, Darshan.patel@hsc.utah.edu
Adam Spivak, MD, adam.spivak@hsc.utah.edu
Joseph P. Alukal, MD, jpa2148@cumc.columbia.edu
Xiaoping Zhang, MD, xzhang@hust.edu.cn
Chengliang Xiong, MD, clxiong951@sina.com
Philip S Li, MD, psli@med.cornell.edu
Honggang Li, MD, lhgyx@hotmail.com
James M. Hotaling, MD, MS, Jim.hotaling@hsc.utah.edu

Corresponding Authors:

James M. Hotaling, MD, MS, FECSM
Associate Professor of Surgery (Urology)
Division of Urology, Department of Surgery
University of Utah School of Medicine
30 N 1900 E, Rm # 3B420
Salt Lake City, UT 84132
Phone: 801-213-2700
Fax 801-585-2891
Email: jim.hotaling@hsc.utah.edu

Abstract Word Count: 195

Word Count: 1,675

Key Words:

COVID-19; Angiotensin Converting Enzyme 2; coronavirus; semen; infertility, male;

Journal Pre-proof

Capsule: SARS-CoV-2 was not detected in semen of COVID-19 patients. *ACE2*-mediated viral entry of SARS-CoV-2 into target host cells is unlikely within the human testicle based on *ACE2* and *TMPRSS2* expression.

Journal Pre-proof

Abstract:

Objective: To describe detection of SARS-CoV-2 in seminal fluid of patients recovering from COVID-19 and describe the expression profile of *ACE2* and *TMPRSS2* within the testicle.

Design: observational, cross-sectional study

Setting: Tertiary referral center

Patients: Thirty-four adult Chinese males diagnosed with COVID-19 through confirmatory quantitative reverse transcriptase-polymerase chain reaction (qRT-PCR) from pharyngeal swab samples

Intervention: None

Main Outcome Measures: Identification of SARS-CoV-2 on qRT-PCR of single ejaculated semen samples. Semen quality was not assessed. Expression patterns of *ACE2* and *TMPRSS2* in the human testis are explored through previously published single-cell transcriptome datasets.

Results: Six patients (19%) demonstrated scrotal discomfort concerning for viral orchitis around the time of COVID-19 confirmation. SARS-CoV-2 was not detected in semen after a median of 31 days (IQR: 29-36 days) from COVID-19 diagnosis. Single-cell transcriptome analysis demonstrates sparse expression of *ACE2* and *TMPRSS2*, with almost no overlapping gene expression.

Conclusions: SARS-CoV-2 was not detected in the semen of patients recovering from COVID-19 one month after COVID-19 diagnosis. *ACE2*-mediated viral entry of SARS-CoV-2 into target host cells is unlikely to occur within the human testicle based on *ACE2* and *TMPRSS2* expression. The long-term effects of SARS-CoV-2 on male reproductive function remain unknown.

Introduction:

SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2), a novel coronavirus reported in late December 2019 in Wuhan, China, has spread world-wide with over 1,000,000 coronavirus disease 2019 (COVID-19) cases reported (1, 2). New developments in molecular virology and immunobiology of SARS-CoV-2, improve our understanding about COVID-19 prevention, management, and possible long-term effects. Although viral transmission occurs predominantly through respiratory droplets, SARS-CoV-2 has been isolated in blood samples and feces from COVID-19 patients, raising questions about viral shedding in other bodily fluids, including semen, as well as alternative modes of transmission(3). Similar to SARS-CoV 2002, viral entry into target cells by SARS-CoV-2 is likely mediated by the interaction between the viral spike (S) protein and cellular angiotensin-converting enzyme 2 (*ACE2*)(4, 5). *ACE2* is expressed in multiple organ systems including type II alveolar cells of the lungs, intestine, heart, kidney, and the testis(4, 5). Transmembrane Serine Protease 2 (*TMPRSS2*) appears to prime the S protein to enhance *ACE2*-mediated viral entry(6). Interestingly, *TMPRSS2* expression is identified in prostatic epithelial cells, with aberrant expression associated with tumorigenesis(7).

The male reproductive tract and the testicle may be involved after some systemic viral infections (e.g., mumps orchitis). Testicular immune privilege normally protects the immunogenic germ cells from the host response. However, certain viruses are able to cross the blood-testis barrier, enter cells of the male reproductive tract, and elicit an immune response within the testicle (8). Evidence regarding viral seeding and viral entry into cells of the male reproductive tract after SARS-CoV-2 infection is not well understood. Our objectives are to 1) describe detection of SARS-CoV-2 in the semen of patients recovering from COVID-19 and 2) determine the expression profile of *ACE2* and *TMPRSS2* within the human testicle, thereby providing mechanistic insights into viral entry and the early impact on male reproductive function.

Methods:

We identified adult Chinese male patients (range 18-57 years) diagnosed with COVID-19 in Wuhan, China between Jan 26th, 2020 and March 2nd, 2020. Local institutional review board approval was obtained prior to study initiation (Huazhong University of Science and Technology, Wuhan, China). A total of 34 adult male patients were recruited for this study after informed consent and were asked to provide one ejaculated semen sample. The duration of abstinence prior to obtaining the semen sample was not standardized.

Patients were initially diagnosed with COVID-19 based on clinical symptoms (fever, cough, pharyngodynia, and respiratory distress) confirmed with quantitative reverse transcriptase-polymerase chain reaction (qRT-PCR) of pharyngeal swab samples (Anda Gene Ltd, Sun yat-sen University, Guangzhou, China). Patients had one or more of these symptoms at the time of COVID-19 confirmation, but generally demonstrated milder symptoms. SARS-CoV-2 nucleic acid was extracted using the automatic nucleic acid extraction system (3DMed®, Shanghai, China), per manufacturer's instruction. Two target genes, including open reading frame 1ab (*ORF1ab*) and nucleocapsid protein (N), were amplified and tested using real-time RT-PCR (Anda Gene Ltd®, Guangzhou, China). A cycle threshold value less than 37 cycles was defined as a positive test, and a value of 40 cycles or more was defined as a negative test, per recommendation of the Chinese National Institute for Viral Disease Control and Prevention. Samples with an equivocal cycle threshold were retested. Detection of SARS-CoV-2 in semen samples was performed using an identical protocol. Semen sample testing for SARS-CoV-2 was performed at the Huazhong University of Science and Technology. Given safety concerns regarding viral transmission from the semen specimens, comprehensive semen analyses were not performed on the samples.

To investigate the gene expression level of *ACE2* and *TMPRSS2* in different cells of the testes, we examined our prior published single-cell RNA-seq (scRNA-seq) dataset of human testicular cells at the University of Utah (9, 10). The scRNA-seq dataset was analyzed as described in prior work (9, 10). In brief, single cell RNA profiling experiments were performed by loading single testicular cells from testes of three healthy, young adults to 10X Genomics® Chromium platform to generate libraries, which were sequenced using Illumina HiSeq 2500®. The generated fastq files were then de-multiplexed and aligned using Cell Ranger®, which will generate the single cell gene expression matrices. The gene expression matrices were then analyzed using dimension reduction algorithm (t-SNE) and clustering approaches. Cells with similar transcriptomes were placed in proximity to each other with the same color. Known markers were utilized to help decode the major testicular cell type identities, as described in prior work (9, 10). Detailed downstream analysis was performed using customized R script. The expression levels of *ACE2* and *TMPRSS2* were examined and displayed by projecting them onto the t-SNE plot. In addition, co-localization of *ACE2* and *TMPRSS2* were reported, and their correlation was calculated.

Results:

Table 1 presents characteristics of the 34 Chinese men recovering from COVID-19. Median age was 37 years (interquartile range, (IQR): 31-49 years). Median body mass index (BMI) was 25.0 kg/m² (IQR: 23.2-26.9), with 17 men (50%) classified as overweight (BMI >25 kg/m²). Three patients (9%) had a past medical history of hypertension. Baseline treatment of hypertension (for example use of ACE inhibitors or angiotensin receptor blockers) was unknown. Interestingly, 6 patients (19%) had scrotal discomfort around the time of COVID-19 confirmation concerning for viral orchitis, although a comprehensive genitourinary examination was not performed on the entire cohort due to the pandemic. Generally, patients in the cohort demonstrated mild-moderate symptoms of COVID-19 at the time of disease confirmation. The median time from the collection of a semen sample from a confirmatory diagnosis of COVID-19 was 31 days (IQR: 29-36 days). SARS-CoV-2 was not detected in any ejaculated semen sample after a median of 31 days.

Figure 1 displays the expression of *ACE2* and *TMPRSS2* using published single-cell transcriptome profiles of human testicular cells. The dimension reduction approach (t-SNE) was utilized to present the single-cell transcriptome data from human testes (6490 single cells), with each dot representing a single cell (Figure 1A). Cells were colored based on cell identities(10). We examined the expression of *ACE2* and *TMPRSS2* by projecting their expression to the t-SNE plot, observing very low expression of both genes (Figure 1B). Last, we examined co-expression of *ACE2* and *TMPRSS2* in every single testicular cell (Figure 1C); only 4 of 6490 cells displayed expression of both genes (Pearson correlation value = -0.01), suggesting limited overlap expression.

Discussion:

In our study, we did not detect SARS-CoV-2 within the semen of adult Chinese males recovering from COVID-19. Nineteen percent of patients in our cohort had scrotal discomfort around the time of their COVID-19 confirmation, although the significance of this remains unclear. Additionally, in our scRNA-seq dataset of human testicular cells at the University of Utah, *ACE2* and *TMPRSS2* are sparsely expressed in the human testes, with almost no overlapping gene expression.

Testicular immune privilege protects the immunogenic germ cells from a host response and a systemic inflammatory response may alter this environment(8). Many viruses such as Mumps,

human immunodeficiency virus (HIV), human herpes virus, Ebola, and Zika can be detected in human semen after infection and can cause orchitis(11, 12). These viruses, through febrile illness, can have a negative impact on male reproductive function and spermatogenesis (11). Before our analysis, the impact of SARS-CoV-2 on male reproductive function was largely unknown. Previously, Xu et al. reported testicular pathology after autopsy from six males who died from complications of SARS-CoV (13). The authors found widespread destruction of germ cells and sperm, in a background of complex inflammatory infiltrate. Although they were unable to isolate a genomic signature from SARS-CoV itself, they postulated that SARS-CoV caused orchitis and reproductive impairment. *ACE2*, a likely receptor for viral entry by SARS-CoV-2, has been previously localized to human Leydig and Sertoli cells(14). Our single-cell transcriptome data suggests *ACE2* RNA expression occurs at low levels. Therefore, *ACE2*-mediated viral entry of SARS-CoV-2 into target host cells is unlikely to occur within the human testicle. Investigation of alternative biologic mechanisms for alteration of the testicular microenvironment by SARS-CoV-2 is warranted.

There are certain limitations to this study. First, our results are impacted by the small sample size and selection bias, as men with COVID-19 in this study are more likely to have demonstrated milder symptoms. Prior research has suggested that higher viral loads are associated with more severe disease symptoms, and it is plausible that viremia or a certain viral threshold is not achieved to cross the blood-testis barrier(15). Second, given safety concerns regarding viral transmission from the semen specimens, comprehensive semen analyses were not performed. We were only able to obtain a single semen sample for the purposes of this study, with only three patients providing a sample within 14 days of their COVID-19 diagnosis. This limits our ability to provide data on possible early viral shedding in the semen. Third, not all patients in our cohort had a comprehensive genitourinary examination, limiting interpretations regarding the incidence of scrotal findings consistent with orchitis during acute SARS-CoV-2 infection. Fourth, we were unable to assess hormone profiles in our cohort, including total testosterone, luteinizing hormone, and follicle stimulating hormone, which could provide an assessment of testicular function. This preliminary data is unable to characterize the long-term impact of COVID-19 on fertility and testicular endocrine function. Finally, we only examined expression of *ACE2* and *TMPRSS2* at the RNA expression level. We understand that gene expression may not always reflect protein abundance, and cannot explicitly conclude that these proteins are absent. Further study of the distribution and roles of *ACE2* and *TMPRSS2* in regulating SARS-CoV-2 infection in the testes is indicated.

Conclusion:

We did not detect SARS-CoV-2 within semen of adult Chinese males recovering from COVID-19, approximately one month after the initial COVID-19 confirmation. Unfortunately, we cannot definitively rule out the presence of SARS-CoV-2 in the seminal fluid during an acute infection with severe COVID-19 symptoms. Additionally, we did not find evidence of *ACE2* and *TMPRSS2* co-expression in high fidelity RNA datasets from our prior published work at the University of Utah, indicating that SARS-CoV-2 would likely not be able to gain entry to testicular cells through an *ACE2/TMPRSS2*-mediated mechanism. Further research is needed to understand the long-term impact of SARS-CoV-2 on male reproductive function including fertility and testicular endocrine function.

Funding:

This work was supported by the National Nature Science Foundation of China (No.81873854) of Feng Pan; National Natural Science Foundation of China (No.81874091) to Xingyuan Xiao; Covid-19 rapid response call of Huazhong University of Science and Technology (2020kfyXGYJ057) of Honggang Li.

Author Contributions:

Category 1:

- a. Conception and Design – Feng Pan, Jingtao Guo, James M. Hotaling
- b. Acquisition of Data - Feng Pan, Xingyuan Xiao, Jingtao Guo, Yarong Song, Honggang Li
- c. Analysis and Interpretation of Data - Feng Pan, Xingyuan Xiao, Jingtao Guo, Yarong Song, Honggang Li, Darshan P. Patel, Adam Spivak, Joseph P. Alukal, Xiaoping Zhang, Chengliang Xiong, Philip S Li, James M. Hotaling

Category 2:

- a. Drafting the Article – Darshan Patel, Jingtao Guo
- b. Revising it for Intellectual Content - Feng Pan, Xingyuan Xiao, Yarong Song, Jingtao Guo, Honggang Li, Darshan P. Patel, Joseph P. Alukal, Xiaoping Zhang, Chengliang Xiong , Philip S Li, James M. Hotaling

Category 3:

- a. Final Approval of the Completed Article - Feng Pan, Xingyuan Xiao, Yarong Song, Jingtao Guo, Honggang Li, Darshan P. Patel, Joseph P. Alukal, Xiaoping Zhang, Chengliang Xiong , Philip S Li, James M. Hotaling

Acknowledgements: None

Table 1: Individual characteristics for adult males with confirmed COVID-19 providing semen sample for SARS-CoV-2 testing.

Patient	Age (yrs)	BMI (kg/m ²)	Time between COVID-19 diagnosis and semen sample obtained (days)	SARS-CoV-2 Semen Result
1	31	26	33	Negative
2	55	23.5	28	Negative
3	31	29.4	32	Negative
4	20	31.2	31	Negative
5	28	24.6	33	Negative
6	50	25.1	29	Negative
7	35	27	27	Negative
8	54	18.1	25	Negative
9	46	24.9	30	Negative
10	30	24.6	40	Negative
11	30	21.7	40	Negative
12	37	29.2	31	Negative
13	49	24.3	31	Negative
14	50	22.2	30	Negative
15	49	26.4	33	Negative
16	55	26.2	37	Negative
17	33	25.1	27	Negative
18	50	23.1	37	Negative
19	37	22	15	Negative
20	27	29.9	31	Negative
21	47	26.6	30	Negative
22	36	22.8	36	Negative
23	36	24.1	33	Negative
24	32	30.9	14	Negative
25	44	26.6	31	Negative
26	32	28.7	33	Negative
27	26	24.1	9	Negative
28	32	32	36	Negative
29	43	22.2	36	Negative
30	39	35.5	8	Negative
31	54	26	35	Negative
32	18	23	29	Negative
33	39	24.3	56	Negative
34	55	22.9	75	Negative

Abbreviations: COVID-19 - coronavirus disease 2019; SARS-CoV-2- severe acute respiratory syndrome coronavirus 2; BMI – body mass index;

Figure 1: Expression of *ACE2* and *TMPRSS2* in human testicular cells using single-cell RNA-seq dataset

A. Dimension reduction (t-SNE) analysis of single-cell transcriptome data from human testes (n = 6490). Reference: Guo et al(10). Each dot represents a single cell and is colored according to its cluster identity, as indicated on the figure key.

B. Expression patterns of *ACE2* and *TMPRSS2* projected on the t-SNE plot. Red indicates high expression, and gray indicates low or no expression, as shown on the figure key.

C. Scatter plot to show the co-expression of *ACE2* (x-axis) and *TMPRSS2* (y-axis) in human testicular cells.

References:

1. Wu Z, McGoogan JM. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *Jama* 2020.
2. . The Center for Systems Science and Engineering, Johns Hopkins, "Coronavirus COVID-19 Global Cases" (2020); [http://www.arcgis.com/apps/opsdashboard/index.html - /bda7594740fd40299423467b48e9ecf6](http://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6).
3. Wang W, Xu Y, Gao R, Lu R, Han K, Wu G *et al.* Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *Jama* 2020.
4. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA *et al.* Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus. *Nature* 2003;426:450-4.
5. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270-3.
6. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S *et al.* SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020.
7. Chen YW, Lee MS, Lucht A, Chou FP, Huang W, Havighurst TC *et al.* TMPRSS2, a serine protease expressed in the prostate on the apical surface of luminal epithelial cells and released into semen in prostasomes, is misregulated in prostate cancer cells. *The American journal of pathology* 2010;176:2986-96.
8. Zhao S, Zhu W, Xue S, Han D. Testicular defense systems: immune privilege and innate immunity. *Cellular & molecular immunology* 2014;11:428-37.
9. Guo J, Nie X, Giebler M, Mlcochova H, Wang Y, Grow EJ *et al.* The Dynamic Transcriptional Cell Atlas of Testis Development during Human Puberty. *Cell stem cell* 2020;26:262-76.e4.
10. Guo J, Grow EJ, Mlcochova H, Maher GJ, Lindskog C, Nie X *et al.* The adult human testis transcriptional cell atlas. *Cell research* 2018;28:1141-57.
11. Liu W, Han R, Wu H, Han D. Viral threat to male fertility. *Andrologia* 2018;50:e13140.
12. Salam AP, Horby PW. The Breadth of Viruses in Human Semen. *Emerging infectious diseases* 2017;23:1922-4.
13. Xu J, Qi L, Chi X, Yang J, Wei X, Gong E *et al.* Orchitis: a complication of severe acute respiratory syndrome (SARS). *Biology of reproduction* 2006;74:410-6.
14. Douglas GC, O'Bryan MK, Hedger MP, Lee DK, Yarski MA, Smith AI *et al.* The novel angiotensin-converting enzyme (ACE) homolog, ACE2, is selectively expressed by adult Leydig cells of the testis. *Endocrinology* 2004;145:4703-11.
15. Liu Y, Yan LM, Wan L, Xiang TX, Le A, Liu JM *et al.* Viral dynamics in mild and severe cases of COVID-19. *The Lancet Infectious diseases* 2020.

