


Rare sugars and their health effects in humans: a systematic review and narrative synthesis of the evidence from human trials

Amna Ahmed , Tauseef A. Khan, D. Dan Ramdath, Cyril W.C. Kendall, and John L. Sievenpiper

Context: Rare sugars are monosaccharides and disaccharides (found in small quantities in nature) that have slight differences in their chemical structure compared with traditional sugars. Little is known about their unique physiological and cardiometabolic effects in humans. **Objective:** The objective of this study was to conduct a systematic review and synthesis of controlled intervention studies of rare sugars in humans, using PRISMA guidelines. **Data Sources:** MEDLINE and EMBASE were searched through October 1, 2020. Studies included both post-prandial (acute) and longer-term (≥ 1 week duration) human feeding studies that examined the effect of rare sugars (including allulose, arabinose, tagatose, trehalose, and isomaltulose) on cardiometabolic and physiological risk factors. **Data extraction:** In all, 50 studies in humans focusing on the 5 selected rare sugars were found. A narrative synthesis of the selected literature was conducted, without formal quality assessment or quantitative synthesis. **Data synthesis:** The narrative summary included the food source of each rare sugar, its effect in humans, and the possible mechanism of effect. Overall, these rare sugars were found to offer both short- and long-term benefits for glycemic control and weight loss, with effects differing between healthy individuals, overweight/obese individuals, and those with type 2 diabetes. Most studies were of small size and there was a lack of large randomized controlled trials that could confirm the beneficial effects of these rare sugars. **Conclusion:** Rare sugars could offer an opportunity for commercialization as an alternative sweetener, especially for those who are at high cardiometabolic risk. **Systematic Review Registration:** OSF registration no. 10.17605/OSF.IO/FW43D.

Affiliation: A. Ahmed, T.A. Khan, C.W.C. Kendall, and J.L. Sievenpiper are with the Department of Nutritional Sciences, Temerty Faculty of Medicine, University of Toronto, Toronto, ON, Canada. A. Ahmed, T.A. Khan, C.W.C. Kendall, and J.L. Sievenpiper are with the Toronto 3D Knowledge Synthesis and Clinical Trials Unit, Clinical Nutrition and Risk Factor Modification Centre, St. Michael's Hospital, Toronto, ON, Canada. D.D. Ramdath is with the Guelph Research & Development Centre, Agriculture and Agri-Food Canada (AAFC), Guelph, Canada. C.W.C. Kendall is with the College of Pharmacy and Nutrition, University of Saskatchewan, Saskatoon, SK, Canada. J.L. Sievenpiper is with the Department of Medicine, Temerty Faculty of Medicine, University of Toronto, Toronto, ON, Canada; Division of Endocrinology and Metabolism, Department of Medicine St. Michael's Hospital, Toronto, ON, Canada; and the Li Ka Shing Knowledge Institute, St. Michael's Hospital, Toronto, ON, Canada.

Correspondence: J.L. Sievenpiper, St. Michael's Hospital, 6138-61 Queen St East, Toronto, ON M5C 2T2, Canada. E-mail: John.sievenpiper@utoronto.ca.

Key words: cardiometabolic health, rare sugars, review.

©The Author(s) 2021. Published by Oxford University Press.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

INTRODUCTION

As rates of obesity and type 2 diabetes continue to rise globally, the role of excess sugars in the diet has become a focus of intense concern.¹ Most of the attention has centered on the adverse health effects of the common sugars – fructose, sucrose, and high-fructose corn syrup (HFCS).² Rare sugars, defined as “monosaccharides and their derivatives that are present in limited quantities in nature”, have received comparatively far less attention.³ These sugars, which can be found in small amounts in a variety of food sources (including honey, certain fruits and vegetables, and grains), may present as unique alternative sweeteners with both caloric and metabolic benefits.^{2,3} Over 40 different types of rare sugars have been identified as having subtle differences in their chemical structure compared with traditional sugars.² Consumption of rare sugars as a sweetener alternative has demonstrated several beneficial physiologic and cardiometabolic effects, including improved glycemic response and weight loss in *in vitro* and animal models. Whether these findings translate to humans and have clinical relevance is unclear.^{4,5} However, evidence of the health effects of rare sugars in humans has begun to accrue for a number of rare sugars, including allulose (psicose), tagatose, isomaltulose (palatinose), L-arabinose, and trehalose. The aim of this review was to provide a systematic summary of the current literature on these rare sugars regarding their physiological and cardiometabolic effects in humans, discuss the possible mechanism for their effects, and highlight their food sources, while also identifying current gaps in the literature on rare sugars.

METHODS

The study followed a systematic search and narrative review methodology.⁶ A systematic search was conducted according to the Cochrane Handbook for Systematic Reviews of Interventions⁷ and reported in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines.⁸ The systematic search was followed by a narrative synthesis of the selected literature, without formal quality assessment or quantitative synthesis. The study protocol was registered as an OSF Registration (osf.io/fw43d) under the following identification number: 10.17605/OSF.IO/FW43D.

Search and selection

MEDLINE and EMBASE were searched through October 1, 2020 for eligible trials. Electronic searches were supplemented with manual searches of references

Table 1 PICOS criteria for inclusion of studies

Parameter	Criterion
Population	Adult humans from all health backgrounds
Intervention	Rare sugars (allulose, L-arabinose, D-tagatose, trehalose, or isomaltulose)
Comparison	Common sugars (sucrose, glucose, maltose, or fructose) Another rare sugar No sugar
Outcomes	Cardiometabolic and physiological risk factors
Study design	Acute (post-prandial studies) trials Longer-term (≥ 1 week duration) trials <i>Included randomized, non-randomized, and uncontrolled human feeding trials</i>

from included studies. Appendix A shows the detailed search strategy. Studies included were randomized, non-randomized, and uncontrolled human feeding trials that examined rare sugars (including allulose, L-arabinose, D-tagatose, trehalose, or isomaltulose) and reported on cardiometabolic and physiological risk factors. Both post-prandial (acute) studies and longer-term (≥ 1 week duration) studies were included. The PICOS (population, intervention, outcome, study design) criteria are provided in Table 1. In all, 882 records were identified through searching MEDLINE and EMBASE. Following removal of duplicates, and after completing a full-text review of the studies identified, 50 studies were eligible for inclusion in this review.

Allulose

Table 2^{9–39} gives the study characteristics for all the included studies on rare sugars. A total of 5 acute and 7 longer-term human studies reported results for allulose and cardiometabolic risk factors. Table 3^{2,37,40–55} gives the chemical and physiological characteristics of rare sugars. Allulose is a monosaccharide found in small amounts in maple syrup, dried fruit, and brown sugar. It is a C-3 epimer of fructose that has about two-thirds of the sweetness of sucrose but a minimal caloric content (0.2 kcal/g).^{40,56} About 70% of allulose is absorbed in the small intestine into the bloodstream (within 1 h) but is excreted intact in urine (within 24 h), while the other 30% is transported to the large intestine, where it is not fermented and thus is excreted intact (within 48 h).⁵⁷ Acute and longer-term randomized controlled trials have examined the effect of allulose consumption on plasma glucose and insulin release, and weight loss, showing benefit in both healthy populations and in individuals with type 2 diabetes.

Table 4^{3,9–22,25–28,30,32,34–37,40,41,58–61} describes the effects of rare sugars in human studies. Kimura et al, in an acute single-bolus randomized controlled trial with healthy participants, examined the effects of

Table 2 Study characteristics

Study, year	Participants	Setting	Mean age, years (SD or range)	Mean BMI, kg/m ² (SD)	Design	Feeding control	Randomization	Rare sugar dose (g)	Intervention or control	Follow-up	Funding sources	Main findings
<i>Allulose (acute)</i>												
Braunstein et al (2018) ⁶⁰	25 H (13 M, 12 F)	OP, Canada	37 (16)	24.7 (3.5)	Crossover	Supp	Yes	5 10	5 g allulose + 75 g OGTT 10 g allulose + 75 g OGTT 75 g OGTT	120 min	A, I	No effect on plasma iAUC or secondary markers of postprandial blood glucose regulation
Iida et al (2008) ⁵⁹	20 H (11 M, 9 F)	OP, Japan	28.2 (6.3)	20.7 (1.8)	Crossover	Supp	Yes	2.5 5 7.5	2.5 g allulose + 75 g OMTT 5 g allulose + 75 g OMTT 7.5 g allulose + 75 g OMTT 75 g OMTT	120 min	NR	Suppressed glucose and insulin levels in a dose-dependent manner ($P < 0.05$)
Kimura et al (2011) ⁵⁸	13 H (5 M, 8 F)	OP, Japan	35.7 (7.6)	20.9 (2.5)	Crossover	Supp	Yes	5	5 g allulose (in 150 mL water) at 30 min prior to MTT 10 g aspartame (in 150 mL water) at 30 min prior to MTT	240 min	NR	Reduction in plasma glucose and increase in fat energy expenditure following test meals ($P < 0.05$)
<i>Control</i>												
Noronha et al (2018) ¹⁰	24 T2DM (12 M, 12 F)	OP, Canada	66 (5.9)	27 (3.4)	Crossover	Supp	Yes	5 10	5 g allulose + 75 g OGTT 10 g allulose + 75 g OGTT 75 g OGTT	120 min	A, I	Reduction in plasma glucose iAUC at 10 g of allulose ($P < 0.05$)
<i>Allulose (longer term)</i>												
Han et al (2018) ⁵⁶	40 OW/OB (20 M, 20 F)	OP, Korea	(20–40)	27.45 (3.21)	Parallel	Supp	Yes	8	8 g/d of allulose in a 30 mL grapefruit flavored noncarbonated beverage	12 wk	A	Reduction in body fat percentage and fat mass at 8 g/d and 14 g/d of allulose ($P < 0.05$)
<i>Control</i>												
Hayashi et al (2010) ¹⁰	8 H (4 M, 4 F) 9 (4 M, 5 F)	OP, Japan	33.4 (3.5) 34.6 (4)	21.3 (2.2) 21.5 (3)	Parallel	Supp	Yes	15	15 g/d allulose in water 15 g/d glucose in water	12 wk	I	No effect on plasma glucose or insulin levels
Hayashi et al (2014) ³	17 OW/OB (8 M, 9 F)	OP, Japan	41.7 (11.5)	25.6 (2.5)	Parallel	Supp	Yes	1.8	30 g/d of a rare sugar syrup containing 6% allulose 28 g/d of high-fructose corn syrup	12 wk	I	Reduction in body weight, fat mass, and waist circumference with consumption of the rare sugar syrup ($P < 0.01$)
<i>Control</i>												
Tanaka et al (2019) ⁹	17 OW/OB (9 M, 8 F)	OP, Japan	42.4 (10.7)	25.4 (2.5)	Open study	Supp	N/A	15	5 g allulose consumed with each meal None	12 wk	I	Increased body fat percentage with allulose consumption ($P < 0.05$)
<i>L-arabinose (acute)</i>												
Krog-Mikkelsen et al (2011) ¹³	15 H (15 M, 0 F)	OP, Denmark	25 (3.2)	22.8 (2.1)	Crossover	Supp	Yes	1 2	1 g L-arabinose + 75 g sucrose in 300 mL water 2 g L-arabinose + 75 g sucrose in 300 mL water	180 min	I	Reduced glucose and insulin peak with L-arabinose consumption ($P < 0.05$)
<i>Control</i>												

(continued)

Table 2 Continued

Study, year	Participants	Setting	Mean age, years (SD or range)	Mean BMI, kg/m ² (SD)	Design	Feeding control	Randomization	Rare sugar dose (g)	Intervention or control	Follow-up	Funding sources	Main findings
Intervention								3	3 g L-arabinose + 75 g sucrose in 300 mL water			
Control									75 g sucrose in 300 mL water			
Shibanuma et al (2011) ¹¹	21 H (18 M, 3 F)	OP, Japan	NR	NR	Crossover	Supp	No	2	2 g L-arabinose + 40 g sucrose dissolved in 108 g deionized water	120 min	NR	Reduction in blood glucose levels at 120 min following L-arabinose consumption ($P < 0.05$)
Control									40 g sucrose in a 150 g sucrose solution			
Halschou Jensen et al (2018) ³⁸	17 H (17 M, 0 F)	OP, Denmark	22.5 (2.6)	22 (1.22)	Crossover	Supp	Yes	2.5 or 2.9	Breakfast and lunch meals supplemented with 5% arabinose by weight	5 h	I	No change in peak plasma glucose or glucose IAUC
Intervention								4.9 or 5.9	Breakfast and lunch meals supplemented with 10% arabinose by weight			
Control									Breakfast and lunch meals			
<i>L-arabinose (longer term)</i>												
Yang et al (2013) ¹²	30 MetS (20 M, 10 F)	OP, China	49.9 (9.9)	NR	Open study	Supp	N/A	40 or 45	20 g twice daily or 15 g thrice daily of L-arabinose	6 mo	I	Reduction in waist circumference ($P < 0.01$), total cholesterol ($P < 0.05$), and fasting glucose ($P < 0.01$)
Control									None			
<i>D-tagatose (acute)</i>												
Buemann et al (2000) ³⁹	20 H (20 M, 0 F)	OP, Denmark	25.7 (4)	24 (2.2)	Parallel	Supp	NR	29	29 g tagatose added to a continental breakfast	13 h	I	Reduced appetite and intake at dinner ($P < 0.05$)
Control									29 g sucrose added to a continental breakfast			
Kwak et al (2013) ¹⁵	52 H (27 M, 25 F)	OP, Korea	35.8 (10.5)	23.7 (3.9)	Crossover	Supp	Yes	5	5 g tagatose sweetened drink + MTT Placebo sweetened (erythritol + 0.004 g sucralose) drink + MTT	120 min	A	Reduction in post-test meal glucose IAUC ($P < 0.05$)
Intervention												
Control												
Kwak et al (2013) ¹⁵	33 pre-diabetes/T2DM (18 M, 15 F)	OP, Korea	57.2 (9.8)	25 (2.6)	Crossover	Supp	Yes	5	5 g tagatose sweetened drink + MTT Placebo sweetened (erythritol + 0.004 g sucralose) drink + MTT	120 min	A	Reduction in post-test meal glucose IAUC ($P < 0.05$)
Intervention												
Control												
Wu et al (2012) ¹⁴	10 H (7 M, 3 F)	OP, Australia	28.8 (12.6)	25.5 (4.7)	Crossover	Supp	Yes	16	40 g tagatose and isomaltulose mixture dissolved in 400 mL water	240 min	NR	Reduced glucose IAUC, serum insulin levels, and slower gastric emptying following the test meal ($P < 0.05$)
Control									60 g sucralose dissolved in 400 mL water			
<i>D-tagatose (longer term)</i>												
Boesch et al (2001) ¹⁶	12 H (12 M, 0 F)	OP, Switzerland	(21–30)	<25	Crossover	Supp	No	45	15 g tagatose added to 3 meals daily	4 wk	NR	No change in body weight
Intervention									15 g sucrose added to 3 meals daily			
Control												
Buemann et al (1998) ¹⁷	8 H (3 M, 5 F)	OP, Denmark	26.2 (2.8)	NR	Crossover	Supp	Yes	30	30 g/d tagatose given in a slice of cake	2 wk	A, I	No change in body weight
Intervention									30 g/d sucrose given in a slice of cake			
Control												
Donner et al (2010) ¹⁹	8 T2DM (4 M, 4 F)	OP, USA	50.7 (10.9)	36.7 (5.1)	Open study	Supp	N/A	45	15 g tagatose taken with food 3 times/day	12 mo	I	Reduction in body weight ($P < 0.05$) and nonsignificant reduction in glycosylated

(continued)

Table 2 Continued

Study, year	Participants	Setting	Mean age, years (SD or range)	Mean BMI, kg/m ² (SD)	Design	Feeding control	Randomization	Rare sugar dose (g)	Intervention or control	Follow-up	Funding sources	Main findings
Control Ensor et al (2015) ²⁰ Intervention	356 T2DM	OP, India & USA	51.7 (10.4)	28.3	Parallel	Supp	Yes	45	None	40 wk	A, I	hemoglobin with tagatose consumption Reduction in body weight ($P < 0.05$) and nonsignificant reduction in glycosylated hemoglobin with tagatose consumption
Control Saunders et al (1999) ¹⁸ Intervention	8 H (4 M, 4 F)	OP, USA	43.6 (5.1)	NR	Parallel	Supp	Yes	75	15 g tagatose dissolved in 125–250 mL of water 3 times/day 1.5 g Splenda dissolved in 125–250 mL of water 3 times/day	8 wk	NR	No change in blood glucose levels, lipid levels, or uric acid
Control Saunders et al (1999) ¹⁸ Intervention	8 T2DM (4 M, 4 F)	OP, USA	53.8 (11.9)	NR	Parallel	Supp	Yes	75	25 g tagatose added to 3 meals daily 25 g sucrose added to 3 meals daily	8 wk	NR	No change in blood glucose levels, lipid levels, or uric acid
Control <i>Trehalose-acute</i> Maki et al (2009) ²² Intervention	23 OB (23 M, 0 F)	OP, USA	49.8 (10.9)	34.9 (0.7)	Crossover	Supp	Yes	75	25 g tagatose added to 3 meals daily No sugar supplementation	120 min	I	Lower rise in plasma glucose and insulin levels ($P < 0.05$)
Control van Can et al (2012) ²¹ Intervention	10 OW (6 M, 4 F)	OP, Netherlands	56 (8)	30.8 (4.9)	Crossover	Supp	Yes	75	75 g trehalose in a 414 mL beverage 75 g glucose in a 414 mL beverage	3 h	I	Lower rise in plasma glucose ($P < 0.01$) and insulin levels ($P < 0.05$)
Control <i>Trehalose (longer term)</i> Kaplon et al (2016) ²⁵ Intervention	32 H (15 M, 17 F) 15 H (7 M, 8 F)	OP, USA	64 (7.7)	NR	Parallel	Supp	Yes	100	75 g trehalose dissolved in 400 mL water 75 g glucose dissolved in 400 mL water	12 wk	A	No change in body weight, lipid levels, or blood pressure
Control Mizote et al (2016) ²³ Intervention	17 H (8 M, 9 F)	OP, Japan	63 (8.2)	26.4 (2.8)	Parallel	Supp	NR	9.9	100 g trehalose mixed with 355 mL of water daily 100 g maltose mixed with 355 mL of water daily	12 wk	NR	Reduction in fasting plasma glucose levels in individuals who had greater trunk fat with trehalose consumption ($P < 0.05$)
Control Yoshizane et al (2020) ²⁴ Intervention	34 MetS (33 M, 1 F) 17 MetS (17 M, 0 F)	OP, Japan	47.9 (7.7)	26.2 (2.6)	Parallel	Supp	Yes	3.3	3.3 g trehalose added to meals and dissolved in drinks 3 times/day 3.3 g sucrose added to meals and dissolved in drinks 3 times/day	12 wk	NR	Plasma glucose levels 2 h after an OGTT closer to fasting plasma glucose levels with trehalose consumption ($P < 0.05$)
Control <i>Isomaltulose (acute)</i> Ang et al (2014) ²⁶ Intervention	50 H (20 M, 30 F)	OP, Japan	43.7 (8.4)	22.4 (3.3)	Parallel	Supp	Yes	3.3	3.3 g trehalose added to meals and dissolved in drinks 3.3 g sucrose added to meals and dissolved in drinks	240 min	NR	Reduced plasma glucose and insulin levels ($P < 0.05$)
Control Arai et al (2005) ³¹ Intervention	11 T2DM (5 M, 6 F)	OP, Germany	53.7 (8.3)	31.6 (4.3)	Crossover	Supp	Yes	1 g/kg BW 1 g/kg BW of sucrose	1 g/kg BW of isomaltulose 1 g/kg BW of sucrose	240 min	A	Reduction in plasma glucose levels at a second meal following isomaltulose beverage consumption ($P < 0.05$)
Control Henry et al (2017) ²⁸ Intervention	7 H (7 M, 0 F)	OP, Japan	31.6 (1.3)	23 (2.6)	Crossover	Supp	Yes	NR	Beverage containing 55.7% isomaltulose Beverage containing 97.2% dextrin	24 h	A	Lower 24 h glucose iAUC ($P < 0.01$) as well as reduced glucose variability with isomaltulose ($P < 0.001$)
Control Henry et al (2017) ²⁸ Intervention	20 H (20 M, 0 F)	OP, Singapore	23.8 (1.8)	24.4 (3.1)	Crossover	Supp	Yes	NR	Breakfast, lunch, and afternoon snack supplemented with isomaltulose			

(continued)

Table 2 Continued

Study, year	Participants	Setting	Mean age, years (SD or range)	Mean BMI, kg/m ² (SD)	Design	Feeding control	Randomization	Rare sugar dose (g)	Intervention or control	Follow-up	Funding sources	Main findings
Control									Breakfast, lunch, and afternoon snack supplemented with sucrose			
Kendall et al (2018) ²⁶ Intervention Control	77 H (18 M, 59 F)	OP, New Zealand	21.9 (5.6)	23.7 (3.6)	Crossover	Supp	Yes	73.2	Trifle containing 73.2 g isomaltulose Trifle containing 73.2 g sucrose	120 min	A	Reduction in blood glucose levels at 60 min following isomaltulose consumption ($P < 0.001$)
Maeda et al (2013) ²² Intervention Control	10 H (10 M, 0 F)	OP, Japan	46.6 (7.7)	21.1 (1.6)	Parallel	Supp	NR	50	50 g isomaltulose dissolved in 300 mL distilled water	180 min	A	Reduction in postprandial plasma insulin and glucose levels following isomaltulose consumption ($P < 0.05$)
Control									50 g sucrose dissolved in 300 mL distilled water			
Sridonpai et al (2016) ³⁰ Intervention Control	11 TZDM	OP, Thailand	49.6 (5.7)	27.8 (2)	Crossover	Supp	Yes	NR	Breakfast supplemented with isomaltulose	240 min	NR	Nonsignificant reduction in plasma glucose levels 30 min–60 min following isomaltulose consumption
Suklaew et al (2014) ²⁷ Intervention Control	12 OB (12 M, 0 F)	OP, Thailand	25.9 (6.6)	25.7 (0.3)	Crossover	Supp	Yes	40	Breakfast supplemented with sucrose 40 g isomaltulose dissolved in 300 mL beverage + high-fat breakfast 40 g sucrose dissolved in 300 mL beverage + high-fat breakfast	480 min	A	Reduction in blood glucose IAUC following isomaltulose consumption ($P < 0.05$)
Control												
<i>Isomaltulose (longer term)</i> Brunner et al (2012) ³⁶ Intervention Control	101 TZDM (66 M, 35 F) 52 TZDM (34 M, 18 F) 49 TZDM (32 M, 17 F)	OP, Germany	60.6 (7.5) 60.5 (8.7)	29.9 (4.2) 32.3 (4.5)	Parallel	Supp	Yes	50	50 g isomaltulose given in biscuits, toffees, milk drinks, and soft drinks	12 wk	I	No difference in HbA1c levels
Holub et al (2009) ³⁷ Intervention Control	20 hyperlipidemic (8 M, 12 F)	OP, Germany	48.2 (21–61)	32.5	Crossover	Met	Yes	50	50 g sucrose given in sweet foods and drinks	4 wk	I	No difference in HbA1c levels
Lightowler et al (2019) ³³ Intervention Control	50 OW/OB	OP, UK	40.7 (11.7)	29.4 (2.7)	Parallel	Supp	Yes	40	40 g isomaltulose + energy-restricted diet 40 g sucrose + energy-restricted diet	12 wk	I	Reduction in weight ($P < 0.001$) and fat mass ($P < 0.01$) with isomaltulose consumption
Mateo-Gallego et al (2019) ³⁴ Intervention Control	43 prediabetes/ TZDM (27 M, 16 F) 21 prediabetes/ TZDM (12 M, 9 F) 22 prediabetes/ TZDM (15 M, 7 F)	OP, Spain	55.9 (6) 55.7 (8.7)	NR	Parallel	Supp	Yes	16.5	Alcohol-free beer supplemented with 16.5 g isomaltulose Alcohol-free beer supplemented with 5.28 g dextrin	10 wk	A	Reduction in HOMA-IR and insulin levels with isomaltulose consumption ($P < 0.05$)
Okuno et al (2010) ³⁵ Intervention Control	50 H (10 M, 40 F) 25 H (5 M, 20 F) 25 H (5 M, 20 F)	OP, Japan	52.2 (8.6) 53.2 (9.4)	23 (2.9) 22.7 (2.8)	Parallel	Supp	Yes	40	40 g isomaltulose given in sugar sticks, jelly, and drinks 40 g sucrose given in sugar sticks, jelly, and drinks	12 wk	NR	Reduction in HOMA-IR with isomaltulose consumption ($P < 0.01$)

Data represents mean \pm SD, unless otherwise stated. Abbreviations: A, agency; F, female; H, healthy; I, industry; M, male; Met, metabolic; MetS, metabolic syndrome; MTT, meal tolerance test; N/A, not applicable; NR, not reported; OB, obese; OGTT, oral glucose tolerance test; OMTT, oral maltodextrin tolerance test; OP, outpatient; OW, overweight; Supp, supplemented; TZDM, type 2 diabetes mellitus.

Table 3 Chemical and physiological characteristics of rare sugars

Rare sugar	Structure	Caloric content (kcal/g)	Sweetness (compared with sucrose)	Glycemic index	Gut enzymes and metabolic fate
Allulose	Monosaccharide (C-3 epimer of fructose)	0.2 ⁴⁰	70% ⁴¹	ND	Transported via GLUT5 in the enterocyte and further transported using GLUT2 (same as fructose) ⁴⁹
L-arabinose	Monosaccharide (aldopentose)	0 ⁵⁰	~50% ⁵¹	ND	Inhibited sucrose activity ²
D-tagatose	Monosaccharide (C-4 epimer of fructose)	(~1.5–3) ⁴³	92% ⁴²	3 ⁵²	Transported via GLUT5 in the enterocyte, metabolized via glycolytic pathway (same as fructose), ² fermented in the colon ⁵³
Trehalose	Dissaccharide (2 glucose molecules in an α 1,1-glycosidic linkage)	4 ⁴⁴	~50% ⁴⁵	ND	Trehalose broken down by trehalase in the small intestine into 2 glucose molecules, which are then absorbed ⁵⁴
Isomaltulose (palatinose)	Disaccharide (glucose and fructose in an α 1-6 glycosidic bond)	4 ⁴⁶	50% ⁴⁷	32 ⁵²	Completely but slowly digested by isomaltase ³⁷
Kojibiose	Disaccharide (2 glucose molecules connected by an α 1-2 glycosidic bond)	ND	ND	ND	
Sorbose	Monosaccharide (ketose)	ND	70% ⁴⁸	ND	
D-allose	Monosaccharide (C-3 epimer of glucose)	ND	80% ⁴⁸	ND	Downregulated GLUT1 expression ⁵⁵

consumption of 5 g allulose, compared with that of 10 mg of aspartame, administered as preloads, on the postprandial glycemic response to a test meal consisting of rice and hamburger steak. They showed a reduction in plasma glucose at 90 minutes following the test meal.⁵⁸ Furthermore, ingestion of allulose as a preload resulted in an increase in fat energy expenditure (but a decrease in carbohydrate energy expenditure) at 90 minutes in response to the test meal compared with ingestion of the test meal alone, demonstrating a possible weight-loss effect.⁵⁸ Iida et al demonstrated that, in healthy individuals, 5 g and 7.5 g of allulose consumed as preloads prior to 75 g of maltodextrin suppressed glucose levels in a dose-dependent manner compared with consumption of the maltodextrin.⁵⁹ Braunstein et al, however, found no effect of 5 g or 10 g of allulose on the postprandial plasma glucose response to a 75 g oral glucose tolerance test (OGTT) in a healthy population. However, the results did reach statistical significance in sensitivity analyses when the results were analyzed according to the assigned placebo (as opposed to the pooled placebo), and the magnitude of effect (25% reduction) was similar to that seen in the earlier trials by Kimura et al and Iida et al.⁶⁰ Noronha et al showed an effect of the same interventions in individuals with type 2 diabetes. Ingestion of 10 g of allulose together with a 75 g OGTT resulted in both a lower plasma glucose iAUC and plasma glucose absolute mean compared with a control of water, while a dose of 5 g of allulose had a borderline significant effect.⁴⁰ A systematic review and meta-analysis of acute feeding trials in people with and without diabetes showed that a small dose of allulose (<30 g) reduced the postprandial iAUC glucose response to the oral glucose load by 10%, while there was an indication of a nonsignificant improvement in iAUC insulin.⁶² These randomized controlled trials demonstrate that small doses of allulose can lead to modest improvement in the postprandial glycemic response to co-ingested carbohydrate.

Longer-term randomized controlled trials show a benefit of allulose on adiposity and glycemic control, though the effect has not been consistently shown.^{3,41} Hayashi et al compared consumption of a beverage sweetened with a rare sugar syrup (containing 6% allulose) daily for 12 weeks with that of a caloric-equivalent beverage sweetened with HFCS, and showed a reduction in body weight, fat mass, and waist circumference in the rare sugar syrup group in obese individuals.³ In addition to allulose, this rare sugar solution also contained glucose, fructose, mannose, sorbose, and other oligosaccharides, making it difficult to attribute the effect entirely to the allulose content.³ It should be noted, however, that the oligosaccharide content of the rare sugar syrup was similar to that of the HFCS

Table 4 Rare sugars and their effects in human studies

Rare sugar	Health-related effects			Side effects
	Healthy individuals	Obese/overweight individuals	Individuals with type 2 diabetes/ borderline type 2 diabetes	
Allulose	<p>Acute: -reduced plasma glucose post-test meal^{58,59} -no effect on plasma glucose⁶⁰ -increased FEE; decreased CEE⁵⁸ Longer term: -reduced BF⁴¹</p> <p>Acute -reduced insulin and glucose peak post-test meal^{11,13}</p>	<p>Long term: -Reduced BW³ -Reduced fat mass³</p> <p>Longer term -reduced WC¹² -reduced TC¹² -reduced fasting plasma glucose¹²</p>	<p>Acute: -Reduced glucose iAUC⁴⁰ Longer term: -No effect on plasma glucose or insulin¹⁰ -Increased BF⁹</p>	<p>Diarrhea, abdominal pain, distension⁴¹</p>
L-arabinose	<p>Acute -reduced insulin and glucose peak post-test meal^{11,13}</p>	<p>Longer term -reduced WC¹² -reduced TC¹² -reduced fasting plasma glucose¹²</p>		<p>Nausea, abdominal pain, diarrhea^{12,13}</p>
D-tagatose	<p>Acute -appetite suppression⁶¹ -lower glucose iAUC¹⁴ Longer term -no effect on BW^{16,17} -no effect on plasma glucose levels¹⁸ Longer term -no effect on BW²⁵</p>		<p>Acute -reduced glucose iAUC¹⁵ Longer term -reduced BW^{19,20}</p>	<p>Nausea, diarrhea, flatulence, bloating^{61,18-20}</p>
Trehalose	<p>Longer term -no effect on BW²⁵</p>	<p>Acute -reduced rise in plasma glucose and insulin levels^{21,22} (3) post-test meal²⁷</p> <p>Acute -reduced plasma glucose post-test meal²⁷ Longer term -reduced BW³³</p>		<p>Bloating, flatulence, diarrhea^{22,25}</p>
Isomaltulose (palatinose)	<p>Acute -reduced plasma glucose post-test meal^{26,28,32} Longer term -reduced HOMA-IR³⁵</p>		<p>Acute -reduced plasma glucose post-test meal³⁰ Longer term -no effect on BW³⁴ -reduced HOMA-IR³⁴ -no effect on HbA1c levels^{36,37}</p>	<p>Diarrhea, nausea, constipation³⁴</p>

Abbreviations: BF: body fat; BW: body weight; CEE: carbohydrate energy expenditure; FEE: fat energy expenditure; HbA1c: hemoglobin A1c; HOMA-IR: Homeostatic Model Assessment of Insulin Resistance; iAUC: incremental area under the curve; TC: total cholesterol; WC: waist circumference.

Table 5 Rare sugars and their FDA, Health Canada, and EFSA designations

Rare sugar	FDA designation	FDA intended use	Health Canada Designation	EFSA designation
Allulose	GRAS Notice 693	Bakery products, beverages, confectionaries, dairy products, sugar substitute, etc.	ND	N.D.
L-arabinose	GRAS Notice 782	Bakery products, baking mixes, condiments, confectionaries, dairy products, snack foods, etc.	ND	N.D.
D-tagatose	ND	ND	ND	Novel food
Trehalose	GRAS Notice 912	Bakery products, frozen desserts, dairy-based foods and toppings, hard and soft confectionery, etc.	Novel food	Novel food

Abbreviations: EFSA, European Food Safety Authority; FDA, Food and Drug Administration; GRAS, generally recognized as safe; ND not described.

intervention. Han et al assessed the effect of two allulose drinks of low (8 g) and high (14 g) dose compared with a 0.024 g sucralose beverage consumed daily for 12 weeks in healthy participants and found a reduction in body fat percentage and fat mass with both allulose drinks; the high dose additionally reduced total subcutaneous fat.⁴¹ Conversely, Tanaka et al found that 15 g/day for 12 weeks of allulose supplementation led to an increase in body fat percentage in 18 diabetic or borderline diabetic participants.⁹ This study lacked a control, and the change in body fat was ascribed by the authors to the additional calories provided by the allulose and the foods with which it was consumed.⁹

In a longer-term randomized controlled trial examining specifically allulose, Hayashi et al demonstrated that 5 g of allulose (compared with 5 g of glucose) 3 times a day for 12 weeks in 17 borderline diabetic participants resulted in no difference in either plasma glucose or insulin levels.¹⁰ A systematic review and meta-analysis of controlled feeding trials of healthy and overweight/obese patients, assessing the effect of small dose of allulose on glycemic markers, did not demonstrate a benefit on hemoglobin A1c (HbA1c) or fasting insulin, though there was a small beneficial effect on fasting glucose.⁶³ In assessing the effect of allulose on cardiometabolic outcomes, Tanaka et al determined that consumption of either 5 g or 15 g of allulose for 48 weeks led to no changes in total cholesterol or LDL cholesterol in 82 hypercholesteremic males and females.⁶⁴ No side effects were observed. Han et al explored gastrointestinal tolerance to allulose in healthy participants and noted symptoms of severe diarrhea only in doses above 0.5 g/kg body weight. When a dose of 0.5 g/kg body weight of allulose was compared with the same dose of sugar, participants reported increased

abdominal pain, distention, and diarrhea. Doses below this threshold, however, were not associated with an increase in the measured gastrointestinal outcomes, indicating that the average individual could consume roughly up to 0.5 g/kg body weight of allulose in a single dose without side effects.⁴¹

In summary, clinical studies show that both the short- and longer-term effects of allulose ingestion may lead to improvement in glycemic outcomes, with possible downstream benefit on body weight and body fat. It is hypothesized that allulose may competitively inhibit movement of glucose into the portal circulation, sharing the same glucose transporter, thereby reducing absorption of glucose in the small intestine.⁴⁰ Additionally, allulose may also increase hepatic glucose uptake, therefore encouraging glycogen synthesis, reducing glucose output from the liver, and reducing glucose plasma levels.⁴⁰ The beneficial effect on glycemic outcomes could also be due to a “catalytic” effect, whereby the small doses of fructose and its epimers may increase the rate-limiting glucokinase activity, leading to a subsequent increase in hepatic glucose metabolism.⁶⁵ In a recent guidance to the industry, the Food and Drug Administration (FDA) concluded, based upon scientific evidence, that allulose is virtually unmetabolized in the human body and thus allowed manufacturers to use a very low 0.4 calories per gram (kcal/g) for allulose. The FDA also concluded that, while allulose is a carbohydrate, based upon its chemical definition, it can be excluded from the “Total Sugars” and “Added Sugars” in a Nutrition Facts label because it is not metabolized, has almost no caloric value, and does not promote dental caries (see Table 5 for regulatory designations for rare sugars).⁵⁷ Overall, as allulose is generally regarded as safe by the FDA, it could prove to be a viable sweetener alternative to sucrose, given its demonstrated physiological and cardiometabolic properties.⁵⁷

L-arabinose

Results from a total of three acute studies and one longer-term human study on L-arabinose and cardiometabolic risk factors have been reported (Table 2). L-arabinose is a monosaccharide and aldopentose found naturally in certain plant cell walls, including many grains and plant gums. It has half the sweetness of sucrose and has been shown in animals to be less metabolizable compared with glucose. With no caloric value, most of the studies examining consumption of L-arabinose in humans are acute post-prandial studies, and they demonstrate a benefit on glycemic control in healthy individuals. All acute trials examining the effect of L-arabinose in humans were conducted using a randomized controlled crossover design. Krog-Mikkelsen et al showed that a number of doses of L-arabinose reduced insulin and glucose peak in healthy males when given prior to a test meal, compared with sucrose. In a similar study design, Shibnuma et al 2010 also found that, in both males and females, consumption of 2 g of L-arabinose before a 40 g sucrose test beverage led to reduced blood glucose levels at 2 hours compared with a control of water.¹¹ However, Halschou-Jensen et al were unable to confirm this effect and found that a breakfast meal supplemented with L-arabinose resulted in no changes in the peak plasma glucose or glucose iAUC compared with a sucrose-supplemented meal in healthy participants.

Yang et al examined the longer-term effect of L-arabinose supplementation in individuals with metabolic syndrome who consumed 40 g–45 g L-arabinose (dissolved in water) daily for 6 months with no alteration in lifestyle habits.¹² This intervention resulted in a reduction in waist circumference, total cholesterol, and fasting glucose, showing an overall benefit in participants with metabolic syndrome.¹² However, since this study lacked a control arm and participants were all diagnosed with metabolic syndrome, it is difficult to extend these results to a larger population. Regardless, the study results promise a novel approach to reducing cardiometabolic risk factors in persons suffering with metabolic syndrome.

No study has specifically examined the side effects of arabinose consumption, though they may occur: the abovementioned study by Krog-Mikkelsen et al showed that out of 15 participants, one experienced mild nausea after 1 g of arabinose, one experienced mild diarrhea after 2 g of arabinose, and another experienced a severe stomach ache and diarrhea after 2 g of arabinose.¹³ Yang et al also noted that, with doses of either 40 or 45 g daily, 13 out of the 30 participants had mild nausea and diarrhea following treatment.¹² A study that specifically examined the gastrointestinal tolerance of

arabinose would be helpful in determining arabinose's side effects and also the maximum recommended dose.

The mechanism by which L-arabinose affects glucose and insulin release in humans is unknown, but in rodent studies it has been shown to inhibit the brush border enzyme sucrose, which can reduce glucose absorption.² Further high-quality studies in humans will be needed to confirm its acute effects and help us to better understand the long-term effects of regular L-arabinose consumption on cardiometabolic outcomes.

D-tagatose

Table 2 shows the study characteristics of 4 acute and 6 longer-term human studies that have reported results for D-tagatose consumption and cardiometabolic risk factors. D-tagatose, a monosaccharide, is a C-4 epimer of D-fructose that is found primarily in whey milk protein and is 92% as sweet as sucrose.⁴² While it has been used as a low-calorie sweetener alternative in milk and yogurt, there is a debate about its exact calorie content, with values ranging from 1.5 kcal/g to 3 kcal/g.^{42,43} Multiple longer-term studies examining the effects of D-tagatose on body weight and blood glucose show a mild benefit.

Randomized acute controlled trials show a benefit for D-tagatose for both glucose and appetite control. In a crossover study, Wu et al demonstrated that in 10 healthy participants a beverage of 40 g D-tagatose and isomaltulose (palatinose), rather than a sucralose beverage, consumed prior to a test meal led to reduced glucose iAUC and serum insulin levels, and slower gastric emptying following the test meal.¹⁴ Buemann et al, using a parallel study design, determined that giving 29 g of D-tagatose in a breakfast meal resulted in reduced appetite and decreased food intake at dinner on the same day in 19 healthy individuals, thus possibly acting as an appetite suppressant.⁶¹ Similarly, Kwak et al conducted an acute cross-over trial in individuals with prediabetes or newly diagnosed type 2 diabetes who were given 5 g of D-tagatose compared with a combination of sucralose plus erythritol. The post-test meal glucose iAUC was reduced with tagatose compared with the control, indicating a benefit in the plasma glucose response.¹⁵

In longer-term studies in healthy individuals, D-tagatose was equivocal in showing benefit. Buemann et al (2 weeks, 8 individuals) and Boesch et al (4 weeks, 12 individuals) both showed no change in body weight with daily ingestion of 30 g and 45 g of D-tagatose, respectively, in a randomized controlled crossover trial.^{16,17} The direction of effect still indicated a possible benefit; it is possible that the effect is small and can only be shown by a different dose or longer duration. In a

randomized controlled parallel trial, Saunders et al similarly examined the effect of 75 g of D-tagatose compared with sucrose daily for 8 weeks in 8 healthy individuals, but saw no change in any of the measured cardiometabolic outcomes, with included blood glucose levels, lipid levels, and uric acid levels.¹⁸

Compared with healthy individuals, the benefit in patients with type 2 diabetes was clearer. In two studies in patients with type 2 diabetes, both Donner et al (8 participants) and Ensor et al (112 participants) demonstrated that the ingestion of D-tagatose resulted in weight loss in a dose- and time-dependent manner.^{19,20} Specifically, Donner et al showed that 45 g/d of D-tagatose for 12 months led to mean reduction of 3.1 kg in an uncontrolled trial, while Ensor et al confirmed this effect with a mean reduction of 5.1 kg in body weight with 45 g/d of D-tagatose for 12 months in a randomized controlled parallel trial.^{19,20} Both studies also showed a non-significant reduction in HbA1C, indicating a possible benefit for blood glucose control in individuals with type 2 diabetes.^{19,20}

Donner et al also noted that all 8 participants had mild gastrointestinal symptoms (including diarrhea, nausea, and flatulence) during the first 2 weeks of D-tagatose supplementation, but these effects subsided for the remainder of the 6 month trial period.¹⁹ Ensor et al reported mild to moderately severe adverse effects, mostly due to gastrointestinal intolerance, with a 5% withdrawal rate due to adverse effects.²⁰ Boesch et al similarly reported diarrhea-like effects and increased bloating in 7 of the 12 participants during the tagatose phase of the study.¹⁶ This was also seen by Saunders et al, in whose study the majority of participants experienced diarrhea and flatulence.¹⁸ Lastly, in a study looking specifically at gastrointestinal tolerance of D-tagatose, Buemann et al determined that approximately 30 g of D-tagatose resulted in diarrhea in approximately 30% of participants, and nausea in approximately 15% of participants, with all individuals reporting flatulence during the 15 day study period.⁶¹ As such, D-tagatose appears to have poor gastrointestinal tolerance at a range of doses, but the effects subside over time.

D-tagatose is known to inhibit the enzymes sucrase and maltase, resulting in reduced absorption of dietary disaccharides, which in turn can increase satiety, potentially explaining the observation of weight loss in human studies.⁶⁶ D-tagatose also promotes hepatic glycogen synthesis and prevents glycogen breakdown, resulting in an increase in glycogen production, leading to reduced plasma glucose levels.⁶⁶ This mechanism, along with the evidence from the trials discussed, demonstrates that D-tagatose shows promise as an alternative sweetener, especially in individuals with type 2 diabetes.

Trehalose

Table 2 gives the study characteristics for all the included studies on trehalose: 2 acute and 3 longer-term human studies reported results for trehalose and cardiometabolic risk factors. Trehalose, a disaccharide of 2 glucose molecules with an α 1,1-glycosidic linkage, is found in yeast, honey, shrimp, insects (for which it is the primary circulating form of energy), and some plants.⁶⁷ It is half as sweet as sucrose, but has the same caloric content.

Trehalose, in acute randomized controlled crossover studies, has been shown to reduce blood glucose levels.^{44,45} Both van Can et al and Maki et al demonstrated that, in overweight adults, consumption of trehalose prior to a test meal led to a lower plasma glucose and attenuated insulin rise when compared with a glucose control.^{21,22} Longer-term parallel controlled studies also show a benefit, but only in individuals with impaired glucose tolerance. Mizote et al determined that 10 g/d of trehalose compared with sucrose for 12 weeks in individuals with metabolic syndrome resulted in a reduction in the fasting plasma glucose levels, but this was limited to those individuals who had greater trunk fat.²³ Furthermore, when participants were stratified by body weight, individuals on the higher end of body weight also saw a reduction in waist circumference and systolic blood pressure.²³ Yoshizane et al similarly showed that, in healthy individuals, consumption of 3.3 g of trehalose for 12 weeks led to plasma glucose levels 2 hours after an OGTT being closer to fasting plasma glucose levels, compared with consumption of sucrose, indicating a benefit in lowering postprandial glucose levels.²⁴ Conversely, Kaplon et al compared 100 g/d of trehalose with 100 g/d of maltose (2 glucose molecules with an α 1,4-glycosidic linkage) for 12 weeks in healthy individuals and saw no difference in body weight, lipid levels, or blood pressure between the groups, indicating a lack of benefit in a healthy population compared with more common sugars.²⁵ However, short- and long-term glucose control measures (such as blood glucose or HbA1c levels) were not measured in this study.

Side effects of trehalose have not been well reported. Kaplon et al saw mild to moderate gastrointestinal discomfort, including bloating, flatulence, and diarrhea in 4 of its 15 patients, while Maki et al reported no adverse effects.^{21,22,25} As such, future studies should also look at the side effects of trehalose at different doses.

Trehalose is metabolized by the brush border enzyme trehalase, which cleaves the 1,1-glycosidic linkage, leaving 2 glucose molecules.²² Trehalase activity, however, is shown to be slower compared with that of other

disaccharidase enzymes, leading to reduced absorption of trehalose and therefore a blunted glucose and insulin response.²² However, trehalose has considerably fewer clinical trials compared with the other rare sugars discussed, and as such needs more long-term clinical and mechanistic studies to substantiate its use as a low-calorie alternative sweetener.

Isomaltulose (palatinose)

A total of 7 acute and 5 longer-term human studies reported results for isomaltulose and cardiometabolic risk factors, as shown in [Table 2](#). Isomaltulose, a more intensely studied rare sugar also known as palatinose, is a disaccharide of glucose and fructose linked together by an α 1-6 glycosidic bond.⁴⁶ Naturally found in small amounts in honey and cane sugar, isomaltulose has half the sweetness of sucrose.^{46,47} While it does have the same caloric content as regular sugar, isomaltulose has been shown to improve the glycemic response in human studies, thus showing promise as an alternative sweetener.^{46,68}

Acute randomized controlled crossover trials demonstrate a benefit for blood glucose and insulin levels from isomaltulose consumption. In an acute trial with 77 healthy adults, Kendall et al demonstrated that consumption of a trifle containing 72.3 g of isomaltulose, compared with one with the same amount of sucrose, led to a reduction in blood glucose levels at 60 minutes following the test meal, with no difference in mean satiety.²⁶ Suklaew et al showed a reduction in glucose iAUC following a meal supplemented with isomaltulose, compared with sucrose, in 12 obese males.²⁷ In a 24-hour study examining supplementation of isomaltulose against sucrose, Henry et al determined that low-glycemic index meals supplemented with isomaltulose led to a lower 24-h glucose iAUC as well as reduced glucose variability over the study period in 20 healthy adults.²⁸ This effect was also confirmed in individuals with type 2 diabetes by Ang et al, who demonstrated that ingestion of 1 g per kg of body weight of isomaltulose compared with sucrose resulted in reduced plasma glucose and insulin concentrations.²⁹ A similar effect was found by Sridonpai et al in individuals with type 2 diabetes, in which an isomaltulose-based breakfast reduced plasma glucose levels 30 to 60 minutes following consumption, compared with a sucrose-based breakfast.³⁰ This effect was carried forward to the next meal, when a standard lunch was given to both groups: those who had an isomaltulose-based breakfast still demonstrated lower plasma glucose levels following the second meal.³⁰ Arai et al confirmed a second-meal effect in 7 healthy males, with plasma glucose and insulin levels remaining low at lunch, following a test breakfast

containing isomaltulose.³¹ Finally, Maeda et al demonstrated that ingestion of 50 g of isomaltulose, compared with 50 g of sucrose, resulted in lower postprandial plasma insulin and glucose levels in 10 healthy males in a parallel controlled trial.³² Overall, isomaltulose shows a benefit in lowering plasma glucose levels acutely compared with sucrose in both healthy participants and in patients with type 2 diabetes, and appears to have an additional second-meal effect.

Longer-term randomized controlled parallel studies also demonstrate a beneficial effect of isomaltulose on cardiometabolic outcomes in both healthy participants and in those at high cardiometabolic risk. In obese and overweight individuals, comparing the intake of 40 g/d isomaltulose with that of sucrose in a calorie-restricted diet, Lightowler et al demonstrated weight loss and reduction in fat mass in the isomaltulose group when given for 12 weeks.³³ Mateo-Gallego et al, on the other hand, did not see any additional effect of isomaltulose on weight loss in a population with type 2 diabetes by administering alcohol-free beer with or without isomaltulose plus maltodextrin for 10 weeks.³⁴ There was, however, a reduction in both Homeostatic Model Assessment of Insulin Resistance (HOMA-IR) and insulin levels in the isomaltulose group but not in the regular alcohol-free beer group, indicating a possible benefit for insulin resistance.³⁴ Okuno et al further confirmed the benefit of isomaltulose for insulin response, because they showed that 40 g/d of a 50/50 isomaltulose and sucrose mix compared with 40 g/d of pure sucrose resulted in a significantly reduced HOMA-IR in healthy adults.³⁵ However, 2 studies, both looking at the effect of 50 g of isomaltulose on glycosylated hemoglobin levels compared with sucrose for either 4 or 12 weeks, found that isomaltulose did not lower HbA1c levels in a population with type 2 diabetes, nor did it affect hyperlipidaemia.^{36,37} The authors hypothesize that the dose provided was not enough to determine a true difference between the effects of isomaltulose and sucrose, suggesting further research is needed on the effects of different doses on HbA1c levels.^{36,37}

Few studies saw any significant side effects of isomaltulose consumption, with only Mateo-Gallego et al reporting that a few participants experienced abdominal discomfort, nausea, diarrhea, and constipation.³⁴ To better understand the maximum tolerable dose of isomaltulose, however, future studies should examine gastrointestinal responses in a dose-dependent manner.

While the exact mechanism of how isomaltulose exerts its effect on plasma glucose and insulin levels has yet to be elucidated, Keyhani-Nejad et al determined that ingestion of isomaltulose (compared with sucrose) resulted in reduced gastric inhibitory polypeptide but increased glucagon-like peptide 1, explaining the

improved metabolic profile seen in many studies.⁶⁹ Overall, while in the literature there is a lack of isomaltulose's effect on body weight, there appears to be an improvement in insulin resistance in several studies, and therefore it may be of some benefit to individuals with type 2 diabetes, though more research is warranted.

Less-studied rare sugars

While there are numerous rare sugars that have yet to be studied in detail, there are a few that show potential in nonhuman studies (cell culture or animal studies). These include kojibiose, sorbose, and allulose. Kojibiose, a glucose disaccharide connected by an α 1-2 glycosidic bond, is found in honey in small amounts.⁷⁰ When examined in vitro in conditions mimicking the upper gastro-intestinal tract and small intestine, kojibiose demonstrated resistance to hydrolysis and was only cleaved by α -glycosidases, and then at a very slow rate.⁷⁰ This delayed digestion might explain the reduced absorption of glucose and may confer a benefit in managing blood glucose levels.⁷⁰ In addition, kojibiose has been shown to be a significant substrate for gut microbiota, creating a beneficial short-chain fatty acid profile, which makes kojibiose a prebiotic.^{70,71} However, kojibiose has yet to be studied in clinical trials; thus, these possible benefits can only be hypothesized for humans.

Similarly, sorbose is a keto monosaccharide with structural similarity to fructose and 70% of the sweetness of table sugar.⁴⁸ It has been well studied in animal models: in long-term studies in rats, sorbose consumption for 2 weeks, compared with sucrose, led to decreased food intake and a reduction in body weight.⁷² Furthermore, acute studies in rats have also demonstrated that sorbose, compared with sucrose, resulted in reduced glucose and insulin levels 30 minutes following ingestion, and the authors identified inhibition of sucrase as a possible mechanism for this result.⁷² Currently, research is needed to identify sorbose's food sources, its effects in humans, its mechanism of action, and its potential as an alternative sweetener.

Lastly, D-allulose, a C-3 epimer of glucose, is 80% as sweet as sucrose and, although its exact caloric content is unknown, it is estimated to be very low in calories.⁴⁸ While these properties would make D-allulose ideal as a low-calorie sweetener, its cardiometabolic effects are not well known, with research instead focusing on its anti-cancer and anti-tumor properties.⁷³ Shown to inhibit proliferation of carcinoma cells and display strong antioxidant characteristics, D-allulose shows benefit in overall inflammation and treatment of disease.⁷³ Future clinical trials should, however, also investigate allulose as a replacement for sucrose, and its subsequent

cardiometabolic effects. Overall, the current literature contains very little information on these rare sugars, in particular whether these sugars would be beneficial as alternative sweeteners, thus providing future areas of research on rare sugars.

Conclusion

Rare sugars, specifically allulose, L-arabinose, D-tagatose, trehalose, and isomaltulose, are exciting in that they may become alternative sweeteners that will offer many physiological and cardiometabolic benefits, ranging from weight loss to improving glycemic control and reducing insulin resistance. Many of these rare sugars need high-quality randomized clinical trials in a larger number of participants coming from a greater variety of health backgrounds, to substantiate many of their benefits. Indeed, for allulose, which has been studied fairly extensively, manufacturers can now state a very low caloric content for it and can also exclude it from "Total sugars" and "Added Sugars" on the Nutrition and Supplemental Facts label in the USA, as per FDA guidance.⁵⁷ Further data elucidating the mechanism of the beneficial effects of these rare sugars is also needed. Given that future research may confirm the safety and benefit of these rare sugars for use as alternative sweeteners, commercialization of these rare sugars could be of great value in helping mitigate the risk associated with diseases such as obesity and type 2 diabetes.

Acknowledgments

Author contributions. A.A., T.A.K. and JLS had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. A.A. and T.A.K. developed and executed the search strategy, extracted the data, performed the analysis and interpretation of the data, and wrote the first draft of the manuscript. D.D.R., C.W.C.K., and J.L.S. participated in the analysis and interpretation of data and critically revised the manuscript for important intellectual content. T.A.K. C.W.C.K. and J.L.S. obtained the funding and were responsible for the original concept, design, and supervision of the work. All authors read and approved the final version of the manuscript.

Funding. The ILSI North America Technical Committee on Carbohydrates contributed to the study design during the grant application process. No other funders contributed to the design, and none of the funders had a role in the conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, approval of the manuscript, or decision to publish, or any other aspect of the present

study. **A.A.** is funded by a Toronto 3D MSc Scholarship Award. **T.A.K.** is funded by a Toronto 3D Post-doctoral Fellowship Award. **J.L.S.** is funded by a Diabetes Canada Clinician Scientist award. The sponsors did not have a role in design or conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, approval of the manuscript, or decision to publish, or any other aspect of the present study.

Declaration of interest. **A.A.** declares no relevant competing interests with the present work.

T.A.K. has received research support from the CIHR and an unrestricted travel donation from Bee Maid Honey Ltd. He has also spoken as an invited speaker at a Calorie Control Council annual general meeting for which he received an honorarium.

D.D.R. has received research support from Pulse Canada, the Saskatchewan Pulse Growers Association, and the Ontario Bean Growers Association. He has no other conflict of interest to declare.

C.W.C.K. has received grants or research support from the Advanced Food Materials Network, Agriculture and Agri-Foods Canada (AAFC), the Almond Board of California, the Peanut Institute, Barilla, the Canadian Institutes of Health Research (CIHR), the Canola Council of Canada, the International Nut and Dried Fruit Council, the International Tree Nut Council Research and Education Foundation, Loblaw Brands Ltd, Pulse Canada, and Unilever. He has received in-kind research support from the Almond Board of California, the American Peanut Council, Barilla, the California Walnut Commission, Kellogg Canada, Loblaw Companies, Nutrartis, Quaker (PepsiCo), Primo, Unico, Unilever, and WhiteWave Foods/Danone. He has received travel support and/or honoraria from the American Peanut Council, Barilla, the California Walnut Commission, the Canola Council of Canada, General Mills, the International Nut and Dried Fruit Council, the International Pasta Organization, Lantmannen, Loblaw Brands Ltd, the Nutrition Foundation of Italy, Oldways Preservation Trust, Paramount Farms, the Peanut Institute, Pulse Canada, Sun-Maid, Tate & Lyle, Unilever, and White Wave Foods/Danone. He has served on the scientific advisory board for the International Tree Nut Council, the International Pasta Organization, the McCormick Science Institute, and Oldways Preservation Trust. He is a member of the International Carbohydrate Quality Consortium (ICQC), an Executive Board Member of

the Diabetes and Nutrition Study Group (DNSG) of the European Association for the Study of Diabetes (EASD), on the Clinical Practice Guidelines Expert Committee for Nutrition Therapy of the EASD and a Director of the Toronto 3D Knowledge Synthesis and Clinical Trials foundation.

J.L.S. has received research support from the Canadian Foundation for Innovation, Ontario Research Fund, Province of Ontario Ministry of Research and Innovation and Science, Canadian Institutes of Health Research (CIHR), Diabetes Canada, PSI Foundation, Banting and Best Diabetes Centre (BBDC), American Society for Nutrition (ASN), INC International Nut and Dried Fruit Council Foundation, National Dried Fruit Trade Association, National Honey Board (the U.S. Department of Agriculture [USDA] honey “Checkoff” program), International Life Sciences Institute (ILSI), Pulse Canada, Quaker Oats Center of Excellence, The United Soybean Board (the USDA soy “Checkoff” program), The Tate and Lyle Nutritional Research Fund at the University of Toronto, The Glycemic Control and Cardiovascular Disease in Type 2 Diabetes Fund at the University of Toronto (a fund established by the Alberta Pulse Growers), and The Nutrition Trialists Fund at the University of Toronto (a fund established by an inaugural donation from the Calorie Control Council). He has received in-kind food donations to support a randomized controlled trial from the Almond Board of California, California Walnut Commission, Peanut Institute, Barilla, Unilever/Upfield, Unico/Primo, Loblaw Companies, Quaker, Kellogg Canada, WhiteWave Foods/Danone, and Nutrartis. He has received travel support, speaker fees and/or honoraria from Diabetes Canada, Dairy Farmers of Canada, FoodMinds LLC, International Sweeteners Association, Nestlé, Pulse Canada, Canadian Society for Endocrinology and Metabolism (CSEM), GI Foundation, Abbott, General Mills, Biofortis, ASN, Northern Ontario School of Medicine, INC Nutrition Research & Education Foundation, European Food Safety Authority (EFSA), Comité Européen des Fabricants de Sucre (CEFS), Nutrition Communications, International Food Information Council (IFIC), Calorie Control Council, and Physicians Committee for Responsible Medicine. He has or has had ad hoc consulting arrangements with Perkins Coie LLP, Tate & Lyle, Wirtschaftliche Vereinigung Zucker e.V., Danone, and Inquis Clinical Research. He is a member of the European Fruit Juice Association Scientific Expert Panel and former member of the Soy Nutrition Institute (SNI) Scientific Advisory Committee. He is on the Clinical Practice Guidelines Expert Committees of Diabetes Canada, European

Association for the study of Diabetes (EASD), Canadian Cardiovascular Society (CCS), and Obesity Canada/Canadian Association of Bariatric Physicians and Surgeons. He serves or has served as an unpaid scientific advisor for the Food, Nutrition, and Safety Program (FNSP) and the Technical Committee on Carbohydrates of ILSI North America. He is a member of the International Carbohydrate Quality Consortium (ICQC), Executive Board Member of the Diabetes and Nutrition Study Group (DNSG) of the EASD, and Director of the Toronto 3D Knowledge Synthesis and Clinical Trials foundation. His wife is an employee of AB InBev.

Supporting Information

The following Supporting Information is available through the online version of this article at the publisher's website.

Table S1 Search term strategy to identify the effects of rare sugars in human studies

Figure S1 Flow of the literature

REFERENCES

- Mitchell NS, Catenacci VA, Wyatt HR, et al. Obesity: overview of an epidemic. *Psychiatr Clin North Am.* 2011;34:717–732.
- Van Laar ADE, Grootaert C, Van Camp J. Rare mono- and disaccharides as healthy alternative for traditional sugars and sweeteners? *Crit Rev Food Sci Nutr.* 2021;61:713–741.
- Hayashi N, Yamada T, Takamine S, et al. Weight reducing effect and safety evaluation of rare sugar syrup by a randomized double-blind, parallel-group study in human. *J Funct Foods.* 2014;11:152–159.
- Kita K, Furuse M, Yang SI, et al. Influence of dietary sorbose on lipogenesis in gold thioglucose-injected obese mice. *Int J Biochem.* 1992;24:249–253.
- Moisés Laparra J, Díez-Munición M, Javier Moreno F, et al. Kojibiose ameliorates arachidic acid-induced metabolic alterations in hyperglycaemic rats. *Br J Nutr.* 2015;114:1395–1402.
- Grant MJ, Booth A. A typology of reviews: an analysis of 14 review types and associated methodologies. *Health Info Libr J.* 2009;26:91–108.
- Higgins JPT, Thomas J, Chandler J, et al. (eds). *Cochrane Handbook for Systematic Reviews of Interventions version 6.0 (updated July 2019)*. Available at www.training.cochrane.org/handbook. Accessed November 2, 2020.
- Moher D, Liberati A, Tetzlaff J, et al.; for the PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *BMJ.* 2009;339:b2535.
- Tanaka M, Hayashi N, Iida T. Safety evaluation of 12-week continuous ingestion of D-allulose in borderline diabetes and type 2 diabetes. *Fundam Toxicol Sci.* 2019;6:225–234.
- Hayashi N, Iida T, Yamada T, et al. Study on the postprandial blood glucose suppression effect of D-psicose in borderline diabetes and the safety of long-term ingestion by normal human subjects. *Biosci Biotechnol Biochem.* 2010;74:510–519.
- Shibanuma K, Degawa Y, Houda K. Determination of the transient period of the EIS complex and investigation of the suppression of blood glucose levels by L-arabinose in healthy adults. *Eur J Nutr.* 2011;50:447–453.
- Yang Z, Li D, Jiang H, et al. The effects of consumption L-arabinose on metabolic syndrome in humans. *J Pharm Nutr Sci.* 2013;3:116–126.
- Krog-Mikkelsen I, Hels O, Tetens I, et al. The effects of L-arabinose on intestinal sucrose activity: dose-response studies *in vitro* and in humans. *Am J Clin Nutr.* 2011;94:472–478.
- Wu T, Zhao BR, Bound MJ, et al. Effects of different sweet preloads on incretin hormone secretion, gastric emptying, and postprandial glycemia in healthy humans. *Am J Clin Nutr.* 2012;95:78–83.
- Kwak JH, Kim MS, Lee JH, et al. Beneficial effect of tagatose consumption on postprandial hyperglycemia in Koreans: a double-blind crossover designed study. *Food Funct.* 2013;4:1223–1228.
- Boesch C, Ith M, Jung B, et al. Effect of oral D-tagatose on liver volume and hepatic glycogen accumulation in healthy male volunteers. *Regul Toxicol Pharmacol.* 2001;33:257–267.
- Buemann B, Toubro S, Astrup A. D-tagatose, a stereoisomer of D-fructose, increases hydrogen production in humans without affecting 24-hour energy expenditure or respiratory exchange ratio. *J Nutr.* 1998;128:1481–1486.
- Saunders JP, Donner TW, Sadler JH, et al. Effects of acute and repeated oral doses of D-tagatose on plasma uric acid in normal and diabetic humans. *Regul Toxicol Pharmacol.* 1999;29:557–565.
- Donner TW, Magder LS, Zarbalian K. Dietary supplementation with D-tagatose in subjects with type 2 diabetes leads to weight loss and raises high-density lipoprotein cholesterol. *Nutr Res.* 2010;30:801–806.
- Ensr M, Williams J, Smith R, et al. Effects of three low-doses of D-tagatose on glycemic control over six months in subjects with mild type 2 diabetes mellitus under control with diet and exercise. *J Endocrinol Diabetes Obes.* 2014;2:1057.
- van Can JGP, van Loon LJC, Brouns F, et al. Reduced glycaemic and insulinaemic responses following trehalose and isomaltulose ingestion: implications for postprandial substrate use in impaired glucose-tolerant subjects. *Br J Nutr.* 2012;108:1210–1217.
- Maki KC, Kanter M, Rains TM, et al. Acute effects of low insulinemic sweeteners on postprandial insulin and glucose concentrations in obese men. *Int J Food Sci Nutr.* 2009;60:48–55.
- Mizote A, Yamada M, Yoshizane C, et al. Daily intake of trehalose is effective in the prevention of lifestyle-related diseases in individuals with risk factors for metabolic syndrome. *J Nutr Sci Vitaminol (Tokyo).* 2016;62:380–387.
- Yoshizane C, Mizote A, Arai C, et al. Daily consumption of one teaspoon of trehalose can help maintain glucose homeostasis: a double-blind, randomized controlled trial conducted in healthy volunteers. *Nutr J.* 2020;19:68.
- Kaplon RE, Hill SD, Bispham NZ, et al. Oral trehalose supplementation improves resistance artery endothelial function in healthy middle-aged and older adults. *Aging (Albany NY).* 2016;8:1167–1183.
- Kendall F, Marchand O, Haszard J, et al. The comparative effect on satiety and subsequent energy intake of ingesting sucrose or isomaltulose sweetened trifle: a randomized crossover trial. *Nutrients.* 2018;10:1504.
- Suklaew P, Suraphad P, Adisakwattana S, et al. The effects of isomaltulose-based beverage on postprandial plasma glucose and lipid profiles in obese men. *J Sci Food Agric.* 2015;1:36–39.
- Henry C, Kaur B, Quek R, et al. A low glycaemic index diet incorporating isomaltulose is associated with lower glycaemic response and variability, and promotes fat oxidation in Asians. *Nutrients.* 2017;9:473.
- Ang M, Linn T. Comparison of the effects of slowly and rapidly absorbed carbohydrates on postprandial glucose metabolism in type 2 diabetes mellitus patients: a randomized trial. *Am J Clin Nutr.* 2014;100:1059–1068.
- Sridonpai P, Komindr S, Kriengsinoyos W. Impact of isomaltulose and sucrose based breakfasts on postprandial substrate oxidation and glycemic/insulinemic changes in type-2 diabetes mellitus subjects. *J Med Assoc Thai Chotmaihet Thangphaet.* 2016;99:282–289.
- Arai H, Mizuno A, Sakuma M, et al. Effects of a palatinose-based liquid diet (Inslow) on glycaemic control and the second-meal effect in healthy men. *Metabolism.* 2007;56:115–121.
- Maeda A, Miyagawa J, Miuchi M, et al. Effects of the naturally-occurring disaccharides, palatinose and sucrose, on incretin secretion in healthy non-obese subjects. *J Diabetes Investig.* 2013;4:281–286.
- Lightowler S, Theis H. Changes in weight and substrate oxidation in overweight adults following isomaltulose intake during a 12-week weight loss intervention: a randomized, double-blind controlled trial. *Nutrients.* 2019;11:2367.
- Mateo-Gallego R, Pérez-Calahorra S, Lamiquiz-Moneo I, et al. Effect of an alcohol-free beer enriched with isomaltulose and a resistant dextrin on insulin resistance in diabetic patients with overweight or obesity. *Clin Nutr.* 2020;39:475–483.
- Okuno M, Kim M-K, Mizu M, et al. Palatinose-blended sugar compared with sucrose: different effects on insulin sensitivity after 12 weeks supplementation in sedentary adults. *Int J Food Sci Nutr.* 2010;61:643–651.
- Brunner S, Holub I, Theis S, et al. Metabolic effects of replacing sucrose by isomaltulose in subjects with type 2 diabetes: a randomized double-blind trial. *Diabetes Care.* 2012;35:1249–1251.
- Holub I, Gostner A, Theis S, et al. Novel findings on the metabolic effects of the low glycaemic carbohydrate isomaltulose (Palatinose™). *Br J Nutr.* 2010;103:1730–1737.
- Halschou-Jensen K, Bach Knudsen KE, Nielsen S, et al. A mixed diet supplemented with L-arabinose does not alter glycaemic or insulinaemic responses in healthy human subjects. *Br J Nutr.* 2015;113:82–88.
- Buemann B, Toubro S, Raben A, et al. The acute effect of D-tagatose on food intake in human subjects. *Br J Nutr.* 2000;84:227–231.
- Noronha JC, Braunstein CR, Glenn AJ, et al. The effect of small doses of fructose and allulose on postprandial glucose metabolism in type 2 diabetes: a double-blind, randomized, controlled, acute feeding, equivalence trial. *Diabetes Obes Metab.* 2018;20:2361–2370.

41. Han Y, Choi B, Kim S, et al. Gastrointestinal tolerance of D-allulose in healthy and young adults. A non-randomized controlled trial. *Nutrients*. 2018;10:2010.
42. Xu Z, Li S, Feng X, et al. L-arabinose isomerase and its use for biotechnological production of rare sugars. *Appl Microbiol Biotechnol*. 2014;98:8869–8878.
43. Turck D, Bresson J, Burlingame B, et al. Scientific Opinion on the energy conversion factor of D-tagatose for labelling purposes. *EFSA J*. 2016;14:e04630.
44. Government of Canada. Archived – novel food information. Available at: <https://www.canada.ca/en/health-canada/services/food-nutrition/genetically-modified-foods-other-novel-foods/approved-products/trehalose.html>. Accessed September 24, 2020.
45. Chattopadhyay S, Raychaudhuri U, Chakraborty R. Artificial sweeteners – a review. *J Food Sci Technol*. 2014;51:611–621.
46. Maresch CC, Pety SF, Theis S, et al. Low glycemic index prototype isomaltulose—update of clinical trials. *Nutrients*. 2017;9:381.
47. Lina BAR, Jonker D, Koziarnowski G. Isomaltulose (Palatinose®): a review of biological and toxicological studies. *Food Chem Toxicol*. 2002;40:1375–1381.
48. Mooradian AD, Smith M, Tokuda M. The role of artificial and natural sweeteners in reducing the consumption of table sugar: a narrative review. *Clin Nutr ESPEN*. 2017;18:1–8.
49. Hishiike T, Ogawa M, Hayakawa S, et al. Transepithelial transports of rare sugar D-psicose in human intestine. *J Agric Food Chem*. 2013;61:7381–7386.
50. Food and Drug Administration. Gras notification for L-arabinose (Betawell® Arabinose). Published online May 4, 2018.
51. Fehér C. Novel approaches for biotechnological production and application of L-arabinose. *J Carbohydr Chem*. 2018;37:251–284.
52. University of Sydney. GI database. Available at: <https://www.glycemicindex.com/foodSearch.php>. Accessed September 24, 2020.
53. Vastenavond CM, Bertelsen H, Hansen SJ, et al. Tagatose (D-tagatose). In O'Brien-Nabors L (ed) *Alternative Sweeteners*. 4th edn. Baton Rouge: CRC; 2011;197–222.
54. Richards AB, Krakowka S, Dexter LB, et al. Trehalose: a review of properties, history of use and human tolerance, and results of multiple safety studies. *Food Chem Toxicol*. 2002;40:871–898.
55. Noguchi C, Kamitori K, Hossain A, et al. D-allose inhibits cancer cell growth by reducing GLUT1 expression. *Tohoku J Exp Med*. 2016;238:131–141.
56. Han Y, Kwon E-Y, Yu M, et al. A preliminary study for evaluating the dose-dependent effect of D-allulose for fat mass reduction in adult humans: a randomized, double-blind, placebo-controlled trial. *Nutrients*. 2018;10:160.
57. Food and Drug Industry. Guidance Document — The Declaration of Allulose and Calories from Allulose on Nutrition and Supplement Facts Labels: Guidance for Industry. Office of Nutrition and Food Labeling Center for Food Safety and Applied Nutrition Food and Drug Administration; 2020.
58. Kimura T, Kanasaki A, Hayashi N, et al. d-Allulose enhances postprandial fat oxidation in healthy humans. *Nutrition*. 2017;43–44:16–20.
59. Iida T, Kishimoto Y, Yoshikawa Y, et al. Acute D-psicose administration decreases the glycemic responses to an oral maltodextrin tolerance test in normal adults. *J Nutr Sci Vitaminol (Tokyo)*. 2008;54:511–514.
60. Braunstein C, Noronha J, Glenn A, et al. A double-blind, randomized controlled, acute feeding equivalence trial of small, catalytic doses of fructose and allulose on postprandial blood glucose metabolism in healthy participants: the Fructose and Allulose Catalytic Effects (FACE) trial. *Nutrients*. 2018;10:750.
61. Buemann B, Toubro S, Astrup A. Human gastrointestinal tolerance to D-tagatose. *Regul Toxicol Pharmacol*. 1999;29:571–577.
62. Braunstein CR, Noronha JC, Khan TA, et al. Effect of fructose and its epimers on postprandial carbohydrate metabolism: a systematic review and meta-analysis. *Clin Nutr*. 2020;39:3308–3318.
63. Noronha J, Braunstein C, Blanco Mejia S, et al. The effect of small doses of fructose and its epimers on glycemic control: a systematic review and meta-analysis of controlled feeding trials. *Nutrients*. 2018;10:1805.
64. Tanaka M, Kanasaki A, Hayashi N, et al. Safety and efficacy of a 48-week long-term ingestion of D-allulose in subjects with high LDL cholesterol levels. *Fundam Toxicol Sci*. 2020;7:15–31.
65. Sievenpiper JL, Chiavaroli L, de Souza RJ, et al. 'Catalytic' doses of fructose may benefit glycaemic control without harming cardiometabolic risk factors: a small meta-analysis of randomised controlled feeding trials. *Br J Nutr*. 2012;108:418–423.
66. Espinosa I, Fogelfeld L. Tagatose: from a sweetener to a new diabetic medication? *Expert Opin Investig Drugs*. 2010;19:285–294.
67. Elbein AD, Pan YT, Pastuszak I, et al. New insights on trehalose: a multifunctional molecule. *Glycobiology*. 2003;13:17R–27R.
68. Oizumi T, Daimon M, Jimbu Y, et al. A palatinose-based balanced formula improves glucose tolerance, serum free fatty acid levels and body fat composition. *Tohoku J Exp Med*. 2007;212:91–99.
69. Keyhani-Nejad F, Kemper M, Schueler R, et al. Effects of palatinose and sucrose intake on glucose metabolism and incretin secretion in subjects with type 2 diabetes. *Diabetes Care*. 2016;39:e38–e39.
70. Beerens K, De Winter K, Van de Walle D, et al. Biocatalytic synthesis of the rare sugar kojibiose: process scale-up and application testing. *J Agric Food Chem*. 2017;65:6030–6041.
71. Díez-Municio M, Kolida S, Herrero M, et al. *In vitro* faecal fermentation of novel oligosaccharides enzymatically synthesized using microbial transglycosidases acting on sucrose. *J Funct Foods*. 2016;20:532–544.
72. Oku T, Murata-Takenoshita Y, Yamazaki Y, et al. D-sorbose inhibits disaccharidase activity and demonstrates suppressive action on postprandial blood levels of glucose and insulin in the rat. *Nutr Res*. 2014;34:961–967.
73. Chen Z, Chen J, Zhang W, et al. Recent research on the physiological functions, applications, and biotechnological production of D-allose. *Appl Microbiol Biotechnol*. 2018;102:4269–4278.