

Subaqueous foraging among carnivorous dinosaurs

<https://doi.org/10.1038/s41586-022-04528-0>

Received: 15 July 2021

Accepted: 7 February 2022

 Check for updates

Matteo Fabbri^{1✉}, Guillermo Navalón^{2,3,4✉}, Roger B. J. Benson^{3✉}, Diego Pol⁵, Jingmai O'Connor¹, Bhart-Anjan S. Bhullar⁶, Gregory M. Erickson⁷, Mark A. Norell⁸, Andrew Orkney³, Matthew C. Lamanna⁹, Samir Zouhri¹⁰, Justine Becker¹¹, Amanda Emke^{11,12}, Cristiano Dal Sasso¹³, Gabriele Bindellini^{13,14}, Simone Maganuco^{13,15}, Marco Audatore¹³ & Nizar Ibrahim¹⁶

Secondary aquatic adaptations evolved independently more than 30 times from terrestrial vertebrate ancestors^{1,2}. For decades, non-avian dinosaurs were believed to be an exception to this pattern. Only a few species have been hypothesized to be partly or predominantly aquatic^{3–11}. However, these hypotheses remain controversial^{12,13}, largely owing to the difficulty of identifying unambiguous anatomical adaptations for aquatic habits in extinct animals. Here we demonstrate that the relationship between bone density and aquatic ecologies across extant amniotes provides a reliable inference of aquatic habits in extinct species. We use this approach to evaluate the distribution of aquatic adaptations among non-avian dinosaurs. We find strong support for aquatic habits in spinosaurids, associated with a marked increase in bone density, which precedes the evolution of more conspicuous anatomical modifications, a pattern also observed in other aquatic reptiles and mammals^{14–16}. Spinosaurids are revealed to be aquatic specialists with surprising ecological disparity, including subaqueous foraging behaviour in *Spinosaurus* and *Baryonyx*, and non-diving habits in *Suchomimus*. Adaptation to aquatic environments appeared in spinosaurids during the Early Cretaceous, following their divergence from other tetanuran theropods during the Early Jurassic¹⁷.

Secondary adaptations to aquatic lifestyles, such as wading behaviour (shoreline specialist and/or only partially submerged habit), subaqueous foraging (fully submerged behaviour) and deep diving, evolved multiple times in every major amniote group^{1,2}. Aquatic habits are widespread among extant birds, ranging from subaqueous foragers to waders. Moreover, water-related ecologies have deep evolutionary roots on the avian stem lineage, occurring in some of the earliest (that is, Early Cretaceous) ornithomorphs^{18,19}. Therefore, the scarcity of evidence for aquatic adaptation in non-avian dinosaurs, which comprise the deep evolutionary stem lineage of birds, is striking.

Non-avian dinosaurs are generally hypothesized to have been restricted to terrestrial environments, with only a few proposed exceptions. Suggestions as to why dinosaurs did not evolve aquatic adaptations as frequently as other amniotes include constraints imposed by the musculoskeletal anatomy of the pelvis, hind limb and tail²⁰. However, the discovery of a new skeleton of the predatory dinosaur *Spinosaurus aegyptiacus* has challenged this long-held narrative^{9,11}: the

conical dentition, retracted nostrils, shortened hindlimbs, paddle-like feet and fin-like tail, together with more ambiguous evidence from isotopic analyses^{21,22} are consistent with an aquatic lifestyle. This has sparked a heated debate regarding the degree of ecological specialization in *Spinosaurus*, which has been described as actively pursuing prey in waters¹¹, with alternative proposals suggesting a more terrestrial or ‘wader-heron’ model on the basis of anatomical observations and 3D digital models (for example, in refs. ^{12,13,23}). Aquatic habits have also been suggested for a handful of other dinosaurs based on gut contents (other spinosaurids²⁴ and ornithomimosaurids^{6,8}) or anatomical proxies (halszkaraptorine dromaeosaurids¹⁰, compsognathids⁵ and various ornithischians^{3,4,7}), but remain ambiguous and controversial. This illustrates the inherent challenges of reconstructing ecomorphological relationships in vertebrates²⁵ and the resulting difficulties in inferring ecological traits in extinct species.

Adaptation to aquatic habits, such as subaqueous foraging or deep diving, constitutes a major evolutionary transition, often culminating

¹Nagaunee Integrative Research Centre, Field Museum of Natural History, Chicago, IL, USA. ²Department of Earth Sciences, University of Cambridge, Cambridge, UK. ³Department of Earth Sciences, University of Oxford, Oxford, UK. ⁴Unidad de Paleontología, Departamento de Biología, Universidad Autónoma de Madrid, Madrid, Spain. ⁵CONICET, Museo Paleontológico Egidio Feruglio, Trelew, Argentina. ⁶Department of Earth and Planetary Sciences and Peabody Museum of Natural History, Yale University, New Haven, CT, USA. ⁷Department of Biological Science, Florida State University, Tallahassee, FL, USA. ⁸Division of Vertebrate Paleontology, American Museum of Natural History, New York, NY, USA. ⁹Section of Vertebrate Paleontology, Carnegie Museum of Natural History, Pittsburgh, PA, USA. ¹⁰Department of Geology and Health and Environment Laboratory, Hassan II University of Casablanca, Casablanca, Morocco. ¹¹Department of Biology, University of Detroit Mercy, Detroit, MI, USA. ¹²Department of Physician Assistant Studies, Wayne State University, Detroit, MI, USA. ¹³Sezione di Paleontologia dei Vertebrati, Museo di Storia Naturale di Milano, Milan, Italy. ¹⁴Dipartimento di Scienze della Terra ‘A. Desio’, Università degli Studi di Milano, Milan, Italy. ¹⁵Associazione Paleontologica Paleontologica Italiana, Parma, Italy. ¹⁶School of the Environment, Geography and Geosciences, University of Portsmouth, Portsmouth, UK. ✉e-mail: mfabbri@fieldmuseum.org; gn315@cam.ac.uk; roger.benson@earth.ox.ac.uk

Table 1 | Phylogenetic regressions comparing explanations of bone compactness as a function of size and ecological traits among femora and dorsal ribs

Femora									
Model	AIC	AIC weights	R ²	Lambda	Variable	Coefficient	Standard error	t	P
Compactness ~ subaqueous foraging	-278.27	0.673	0.172	0.919	Intercept	0.63	0.108	5.8585	0.00
					Subaqueous foraging	0.164	0.023	7.0225	0.00
Compactness ~ subaqueous foraging + sustained flight	-275.35	0.156	0.168	0.915	Intercept	0.6315	0.105	6.0075	0.00
					Subaqueous foraging	0.163	0.023	7.039	0.00
					Sustained flight	-0.056	0.026	-2.151	0.0326
Compactness ~ subaqueous foraging + flight	-275.33	0.154	0.168	0.912	Intercept	0.633	0.104	6.089	0.00
					Subaqueous foraging	0.162	0.023	6.945	0.00
					Flight	-0.057	0.027	-2.41	0.03
Dorsal ribs									
Model	AIC	AIC weights	R ²	lambda	Variable	Coefficient	Standard error	t	P
Compactness ~ subaqueous foraging	-164.167	0.638	0.108	0.969	Intercept	0.648	0.061	10.652	0
					Subaqueous foraging	0.154	0.033	4.7195	0
Compactness ~ subaqueous foraging + sustained flight	-161.538	0.171	0.104	0.97	Intercept	0.651	0.06	10.8315	0
					Subaqueous foraging	0.152	0.032	4.735	0
					Sustained flight	-0.055	0.024	-2.2595	0.02545
Compactness ~ subaqueous foraging + flight	-161.421	0.162	0.104	0.968	Intercept	0.655	0.06	10.9355	0
					Subaqueous foraging	0.148	0.032	4.575	0
					Flight	-0.056	0.025	-2.234	0.0271

Explanatory variables are combined in all possible ways in different linear models. Linear models are ordered from best model to worst. Median values from the 100 phylogenies used for the phylogenetic regressions. Models with negligible AIC weights were trimmed down from this table but can be found in Supplementary Tables 3, 4. These analyses are two-sided and no adjustments were applied for multiple comparisons.

into fundamental transformation of the body plan²⁵. Nevertheless, even in groups that exhibit a high degree of aquatic specialization, such as cetaceans and many marine reptiles, this transformation occurred gradually over millions or tens of millions of years. Some extant species, and the early fossil members of even the most specialized aquatic groups, can show relatively subtle skeletal changes^{14–16,26,27}. Many aquatic taxa possess few anatomical indicators of water-related ecology, and instead share numerous traits with land animals (for example, *Hippopotamus* and the earliest cetaceans^{14,15,26,27}). It is therefore plausible that dinosaurs currently considered to have been terrestrial on the basis of anatomical proxies and phylogenetic bracketing might instead represent the early stages of an evolutionary transition towards more specialized aquatic ecologies (for example, early cetaceans) or amphibious animals (for example, *Hippopotamus* and *Tapirus*) that evolved relatively limited anatomical transformations despite spending much of their lives in water.

Because of the difficulty of inferring aquatic habits from skeletal morphology alone, proxies that reveal ecological adaptations in extinct taxa are required. Osteohistological features such as variation in bone density provide one such possibility. Osteosclerosis occurs widely as

an adaptation to aquatic life in extant amniotes^{26–29}, and has been used to infer aquatic ecologies in extinct tetrapods such as crocodyliforms, avialans, marine reptiles and cetaceans^{15,26,27}. Osteosclerosis involves additional deposition of bone mass per volumetric unit leading to the presence of a thick bone cortex with dense trabecular networks infilling the medullary cavity^{2,26–29}. This results in increased body density, facilitating buoyancy control during subaqueous immersion related to either submerged aquatic foraging (for example, in underwater pursuit divers), concealment or refuge^{14–16,26–29}. Although previously used for paleoecological inference, bone density has generally been used on single-clade-specific studies for example, in ref.³⁰), and a phylogenetically broad test is required to validate the use of bone compactness as a proxy for aquatic adaptation in deep time, including in species outside of the extant crown clades such as non-avian dinosaurs.

Here we conduct phylogenetic comparative analyses of bone density data in a broad sample of amniotes and use our findings to assess the extent of aquatic adaptations in non-avian dinosaurs. Our analyses provide evidence that one clade of dinosaurs—Spinosauridae—was ecologically adapted to life in water, representing the first known aquatic radiation among non-avian dinosaurs.




Fig. 1 | Osteohistology and ecological variation among amniotes, including the analysed spinosaurid taxa.

a, Bipedal, land-dwelling archosaurs such as theropods show the presence of an open medullary cavity. This condition is more pronounced in flying archosaurs such as birds. Two osteosclerotic patterns are present among subaqueous foraging animals: (1) increase in thickness of the bone cortex, as observed in crocodylians and penguins, for animals adapted to shallow waters; or (2) substitution of the bony cortex with trabecular networks, usually found in deep divers—for example, ichthyosaurs, mosasaurs and cetaceans. Occupation of the medullary cavity by spongiosa is also observed in quadrupedal, graviportal animals such as sauropods,

ornithischians and large-bodied terrestrial mammals. **b**, Femur and dorsal rib sections and bone density of the holotype of *Baryonyx*, *Suchomimus* and the neotype of *Spinosaurus* used for calculation of bone density in this study. Skeletal reconstructions are based on single individuals (holotype of *Baryonyx* and neotype of *Spinosaurus*), exception made for *Suchomimus* (see Supplementary Information for further details); preserved bones are highlighted in orange. The schematic tree is based on the phylogenetic analyses performed in this study (see Supplementary Information for results and discussion of these analyses). bd, bone density.

We quantify bone density in the femoral diaphysis and proximal region of dorsal ribs of 206 and 174 extant and extinct amniotes, respectively (380 total observations with $n = 83$ overlapping taxa between the two datasets; see Supplementary Dataset and Supplementary Table 1). Our dataset includes novel osteohistological data for non-avian dinosaurs (36 femora and 12 ribs) and Mesozoic stem-avialans (7 femora) (Extended Data Figs. 1–7 visualize novel data used in this study; see Methods, Supplementary Dataset for the list of taxa and bone density values used in this study and Supplementary Table 2 for a list of spinosaurids and their investigated skeletal elements).

We compared alternative explanations of variation in bone density using corrected Akaike information criterion (AIC)-based model comparison of phylogenetic multiple regressions³¹, and evaluated the influence of allometry using the maximum diameter of the femoral diaphysis and proximal region of the dorsal ribs as a size proxy. As ecological adaptations are often reflective of the most demanding biomechanical behaviour (Liem's paradox³²: whereas specialized animals are capable of less functionally demanding behaviours, less specialized taxa often cannot satisfy the requirements linked to functionally challenging habits such as sustained flight or subaqueous foraging), taxa were scored using two categorical explanatory variables that encode the presence of (1) subaqueous foraging (0, unable; 1, able but infrequent;

2, frequent), and (2) flying (0, unable; 1, non-sustained flight; 2, sustained flight) in a comprehensive evolutionary framework. We used two independently varying variables because flight and subaqueous foraging evolved at least partly independently of one another as indicated by the occurrence of both flying and flightless diving birds. Our datasets include extant and extinct taxa with undisputed aquatic adaptations, specifically marine mammals (cetaceans and pinnipeds) and non-archosaurian marine reptiles (ichthyosaurs and sauropterygians, including plesiosaurs and mosasaurs), and aquatic archosaurs such as metriorhynchids, living crocodylians and various clades of subaqueous foraging birds (penguins, auks, loons, grebes and cormorants), in addition to extant and extinct terrestrial and flying archosaurs, lepidosaurs and mammals (see Supplementary Dataset).

The best linear model is 'bone compactness - subaqueous foraging' (state 2: frequent subaqueous foraging) in both datasets (AIC of weight = 0.673 (femur) and 0.638 (rib); Table 1, Supplementary Tables 3–4). This indicates that frequent subaqueous foraging is associated with increased femoral and rib density across amniotes ($P < 0.001$), a relationship that exhibits a strong phylogenetic signal (median $\lambda = 0.97$ (femur) and 0.969 (rib)). Infrequent subaqueous foraging and wading behaviour are not significantly associated with variation in bone density (Table 1, Supplementary Tables 3, 4), consistent with the observation that wading birds that feed in water but rarely

Q11





Fig. 2 | Relationship between midshaft density of femur, diameter and major lifestyle among amniotes including Spinosauridae.

a, Phylogenetically gnostic regressions (PGLS) linear model ($n = 206$ independent observations, $n = 200$ taxa) of bone density as a function of \log_{10} femur-midshaft diameter values for our dataset of amniotes with main lifestyle category as a factor using a randomly drawn phylogeny from the 100 phylogenies generated (same topology, variable branch lengths). Solid lines represent linear fits for the four categories. The dashed line represents linear fit for aquatic taxa without deep diving and graviportal taxa (asterisks inside

Q12

submerge (for example, herons, pelicans, gulls, flamingoes and some ducks) have similar compactness to non-aquatic taxa.

Models that include flight or shaft diameter as additional covariates receive less support from AIC (Table 1, Supplementary Tables 3, 4). This indicates that evidence for an amniote-wide common allometry in bone density, or for association of flight with decreased skeletal density, is weak, and that those effects are secondary to that of aquatic adaptation (see Table 1, Extended Data Figs. 8, 9, Supplementary Tables 3, 4). Nevertheless, negative allometry in bone compactness (reduction of bone density with size increase) is found in flying taxa when analysed separately (volant extant birds, Cretaceous enantiornithines and pterosaurs; Table 1, Supplementary Tables 3, 4). This shows the importance



the dot symbol). **b**, Violin plots depicting distribution of bone density in each category. Large dots represent medians and lines show 95% confidence intervals. The bone densities of the three spinosaurid taxa studied are indicated. **c**, Bone density distribution in a time-calibrated archosaur consensus tree showing an ancestrally osteosclerotic Spinosauridae and rapid radiation of tetanuran clades during the Early Jurassic (Extended Data Fig. 8 shows bone density distribution across amniotes). Eoc., Eocene; Mio., Miocene; Olig., Oligocene; Paleo, Palaeogene; Quat., Quaternary.

of skeletal weight reduction in association with or preceding the origin of active flight^{33,34}: postcranial bones of predatory dinosaurs typically show an open medullary cavity, a trait inherited by birds^{14,34} (Fig. 1, Extended Data Figs. 1–9). Large-bodied terrestrial amniotes have relatively high femoral compactness related to graviportal taxa: trabeculae invade the medullary cavity to support increased weight in graviportal mammals^{15,27,35,36} and sauropod dinosaurs (Fig. 1, Extended Data Figs. 1–7, Supplementary Tables 5, 6). Deep diving animals, such as ichthyosaurs, mosasaurs, living cetaceans and seals, are characterized by lower bone density when compared to shallow-water subaqueous foragers: the compact bone cortex of deep divers is replaced by cancellous bone characterized by extensive trabeculae and vascularization^{2,27} (Fig. 1,




Fig. 3 | Relationship between dorsal ribs density, diameter and major lifestyle among amniotes including Spinosauridae. **a.** PGLS linear model ($n = 174$ taxa) of bone density as a function of \log_{10} rib cross-section diameter values for our dataset of amniotes with main lifestyle category as a factor using a randomly drawn phylogeny from the 100 phylogenies generated (same topology, variable branch lengths). Solid lines represent linear fits for the four

categories. Dashed line represents linear fit for aquatic taxa without pelagic and graviportal taxa (asterisks inside the dot symbol). **b.** Violin plots depicting distribution of bone compactness values in each category. Large dots represent medians and lines show 95% confidence intervals. Bone compactness of *Baryonyx* and *Spinosaurus* are indicated (Extended Data Fig. 9 shows bone compactness distribution across amniotes).

Extended Data Figs. 1–7, Supplementary Tables 5, 6), hypothesized as counteracting compression in deep waters and increases in metabolism¹². High bone density is therefore an excellent indicator for the initial stages of aquatic adaptation, but poorly distinguishes between wading, deep diving, and terrestrial habits. These limitations can be overcome using anatomical observations because deep diving show other transformations of the body plan, such as presence of fins and flippers. Graviportal animals can be distinguished from aquatic species by the presence of columnar limbs, an anatomical trait which is generally missing among subaqueous foragers. Furthermore, graviportal taxa do not affect rib compactness (Extended Data Figs. 1–7, Supplementary Dataset). These analyses therefore demonstrate that bone density is a powerful proxy of shallow subaqueous foraging across amniotes.

We used this relationship to establish quantitative predictions of subaqueous foraging in a range of non-avian dinosaurs, including groups that were previously suggested to be linked to water^{4,6,8–11}, using phylogenetically flexible discriminant analyses with all amniotes in our sample (Methods). We repeated analyses across 100 informal supertrees with varying branch lengths to account for stratigraphic uncertainty. The informal consensus trees include a novel phylogenetic analysis of Tetanurae modified from recently published datasets^{17,37}, including new observations of the *Spinosaurus* neotype (Figs. 1–3, Supplementary Fig. 1; Supplementary Materials). Our analyses include novel osteohistological data for the spinosaurids *Baryonyx*²⁴, *Suchomimus*^{9,38} and *Spinosaurus*^{9,11}, as well as other tetanuran theropods (see Supplementary Materials for ontogenetic assessments of these taxa and other carnosaurs analysed in this study; Fig. 1, Extended Data Fig. 10, Supplementary Figs. 2, 3).

The correct classification rates of our phylogenetically flexible discriminant analyses ranges are 84–85% (femora) and 83–84% (ribs) (Figs. 2, 3, Supplementary Materials, Supplementary Tables 7–10). This increases to 90% in both datasets when excluding graviportal and deep diving taxa (Figs. 2, 3, Supplementary Tables 7–10). Contrary to previous hypotheses, our analyses indicate that Spinosauridae is the only clade of non-avian dinosaurs with unambiguous evidence of subaqueous foraging. Within Spinosauridae, disparate ecomorphologies were found. *Spinosaurus* (median probability for subaqueous

foraging 100% (femur) and 95% (rib)) and *Baryonyx* (median probability for subaqueous foraging 98% (femur) and 96% (rib)) were predicted as subaqueous foragers. By contrast, *Suchomimus* was found as non-subaqueous-forager (median probability for subaqueous foraging 31% (femur)), similar to other terrestrial non-avian dinosaurs (Figs. 1–3, Extended Data Fig. 10, Supplementary Tables 7–10). Considering the similar body size between the skeletally mature specimen of *Suchomimus* (G51) and the neotypic skeletally immature individual of *Spinosaurus*, and that our analytical approach accounts for size variation, these results can be confidently attributed to ecological adaptations, rather than the influence of allometry. This is also supported by the presence of open medullary cavities in postcranial elements of other large bodied, bipedal predatory dinosaurs such as femora of *Tyrannosaurus*, *Tyrannotitan*, *Torvosaurus*, and a large carcharodontosaurid rib, contrasting with the osteosclerotic bones of *Baryonyx* and *Spinosaurus* (Fig. 1, Extended Data Figs. 1–7, 10).

All other investigated non-avian dinosaur clades (ornithomimosaurs^{6,8}, halszkaraptorine dromaeosaurids¹⁰ and ornithopods^{3,4,7}), also show open medullary cavities and a weak or absent probability of subaqueous foraging (see Supplementary Tables 7–10). By contrast, the inference of subaqueous foraging in some spinosaurids is especially convincing because osteosclerosis is observed across multiple skeletal elements in both the holotype of *Baryonyx* (dorsal ribs, scapula, pubis, ischium, femur and fibula) and the neotype of *Spinosaurus* (dorsal ribs, dorsal and caudal neural spines, femur, tibia, fibula and manual phalanx) (Fig. 1, Extended Data Fig. 10, Supplementary Figs. 2, 3), rendering previous biomechanical models inaccurate¹².

Phylogenetic optimization of bone density and the presence of osteosclerosis tentatively suggests that subaqueous foraging, is ancestral for Spinosauridae (Fig. 2c, Extended Data Figs. 8, 9) and that the absence of osteosclerosis in *Suchomimus* results from secondary loss rather than primitive absence. The absence of osteosclerosis does not rule out a dependency on aquatic habitats for predation in *Suchomimus*: anatomical traits are consistent with a largely piscivorous diet, including an elongate snout and conical dentition. One possibility is that *Suchomimus* was a wading predator hunting from riverbanks, as previously hypothesized for other spinosaurids^{12,13,24,38–40}. Different ecological

adaptations (subaqueous foraging and non-diving habits) are recovered between the anatomically similar sister taxa *Baryonyx* and *Suchomimus*, a pattern not unique to Baryonychinae and also observed in other amniote groups, including Phalacrocoracidae (this study) and Hippopotamoidea³⁵. It is possible that environmental factors, such as a sparser distribution of aquatic settings (rivers, lakes)⁴¹ led to less specialized foraging in *Suchomimus*.

Our results suggest the first anatomical adaptations for an aquatic lifestyle appeared in concert with osteosclerosis in spinosaurids. Craniofacial modifications preceded postcranial alterations (Fig. 1). The premaxilla gradually became more elongate, while the external naris diminished in size and migrated posterodorsally^{9,11}, a pattern comparable to the telescoping process observed in the skull evolution of cetaceans¹⁴ and ichthyosaurs¹⁶. The braincase rotated ventrally and the dentition became conical. These modifications are functionally advantageous for a diet based on slippery, aquatic prey⁴². Postcranial modifications linked to subaqueous foraging, such as elongation of the caudal neural spines to form a propulsive structure, have been reported for *Spinosaurus*^{9,11} and the baryonychine *Riparovenator*⁴⁰. Additionally, spinosaurids are characterized by the lowest degree of postcranial pneumatization (restricted to the cervical region and dorsal–sacral neural arches) among large-bodied Cretaceous tetanurans⁴³, consistent with elevated body density and advantageous for buoyancy control. Although reduction in hind limb length and widening of the pes have only been described in *Spinosaurus*^{9,11}, many spinosaurids are only known from fragmentary remains (Fig. 1), limiting our understanding of their skeletal adaptations. Because of their unique anatomy, spinosaurids may have had ecologies with no modern equivalent, limiting direct autecological interpretations based on modern taxa.

We demonstrate that Spinosauridae, a geographically widespread clade of predatory dinosaurs, was ecomorphologically adapted to life in water, but that aquatic adaptation was otherwise absent among non-avian dinosaurs studied so far. Nonetheless, this finding challenges the hypothesis that non-avian dinosaurs were restricted to terrestrial environments. Spinosaurids were part of the rapid radiation of Tetanurae during the late stages of the Early Jurassic¹⁷. Increased diversification appeared in concert with morphological innovation and high rates of homoplasy across tetanurans¹⁷. This ecomorphological radiation may be linked to adaptation to previously under-exploited environments, including multiple independent appearances of aerial capabilities³³. Our study demonstrates that ecomorphological radiations among non-avian dinosaurs also included the invasion of freshwater ecosystems.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41586-022-04528-0>.

- Kelley, N. P. & Pyenson, N. D. Evolutionary innovation and ecology in marine tetrapods from the Triassic to the Anthropocene. *Science* **348**, aaa3716 (2015).
- Gutarra, S. & Rahman, I. A. The locomotion of extinct secondarily aquatic tetrapods. *Biol. Rev.* **97**, 67–98 (2022).
- Owen, R. A description of a portion of the skeleton of the *Cetiosaurus*, a gigantic extinct saurian reptile occurring in the oolitic formations of different portions of England. *Proc. Geol. Soc. Lond.* **3**, 457–462 (1841).
- Cope, E. On the characters of the skull in the Hadrosauridae. *Proc. Nat. Acad. Nat. Sci.* **35**, 97–107 (1883).
- Bidar, A., Demay, L. & Thomel, G. *Compsognathus corallestris*, une nouvelle espèce de dinosaure thérope du Portlandien de Canjuers (Sud-Est de la France). *Annales Muséum d'Histoire Naturelle de Nice* **1**, 9–40 (1972).
- Norell, M. A., Makovicky, P. J. & Currie, P. J. The beaks of ostrich dinosaurs. *Nature* **412**, 873–874 (2001).
- Tereschenko, V. S. Adaptive features of protoceratopoids (Ornithischia: Neoceratopsia). *Paleontol. Journ.* **42**, 273–286 (2008).

- Lee, Y. N. et al. Resolving the long-standing enigmas of a giant ornithomimosaur *Deinocheirus mirificus*. *Nature* **515**, 257–260 (2014).
- Ibrahim, N. et al. Semiaquatic adaptations in a giant predatory dinosaur. *Science* **345**, 1613–1616 (2014).
- Cau, A. et al. Synchrotron scanning reveals amphibious ecomorphology in a new clade of bird-like dinosaurs. *Nature* **552**, 395–399 (2017).
- Ibrahim, N. et al. Tail-propelled aquatic locomotion in a theropod dinosaur. *Nature* **581**, 67–70 (2020).
- Henderson, D. M. A buoyancy, balance and stability challenge to the hypothesis of a semi-aquatic Spinosaurus Stromer, 1915 (Dinosauria: Theropoda). *PeerJ* **6**, e5409 (2018).
- Hone, D. W. E. & Holtz, T. R. Jr Evaluating the ecology of *Spinosaurus*: shoreline generalist or aquatic pursuit specialist? *Palaeontol. Electronica* **24**, a03 (2021).
- Thewissen, J. G., Cooper, L. N., Clementz, M. T., Bajpai, S. & Tiwari, B. N. Whales originated from aquatic artiodactyls in the Eocene epoch of India. *Nature* **450**, 1190–1194 (2007).
- Houssaye, A. Bone histology of aquatic reptiles: what does it tell us about secondary adaptation to an aquatic life? *Biol. J. Linn. Soc.* **108**, 3–21 (2013).
- Motani, R. et al. A basal ichthyosauriform with a short snout from the Lower Triassic of China. *Nature* **517**, 485–488 (2015).
- Rauhut, O. W. & Pol, D. Probable basal allosauroid from the early Middle Jurassic Cañadón Asfalto Formation of Argentina highlights phylogenetic uncertainty in tetanuran theropod dinosaurs. *Sci. Rep.* **9**, 1–9 (2019).
- You, H. L. et al. A nearly modern amphibious bird from the Early Cretaceous of northwestern China. *Science* **312**, 1640–1643 (2006).
- Wilson, L. E. & Chin, K. Comparative osteohistology of *Hesperornis* with reference to pygoscelid penguins: the effects of climate and behaviour on avian bone microstructure. *R. Soc. Open Sci.* **1**, 140245 (2014).
- Gatesy, S. M. & Dial, K. P. Locomotor modules and the evolution of avian flight. *Evolution* **50**, 331–340 (1996).
- Amiot, R. et al. Oxygen isotope evidence for semi-aquatic habits among spinosaurid theropods. *Geology* **38**, 139–142 (2010).
- Hassler, A. et al. Calcium isotopes offer clues on resource partitioning among Cretaceous predatory dinosaurs. *Proc. R. Soc. B* **285**, 20180197 (2018).
- Larramendi, A., Paul, G. S. & Hsu, S. Y. A review and reappraisal of the specific gravities of present and past multicellular organisms, with an emphasis on tetrapods. *Anat. Rec.* **304**, 1833–1888 (2021).
- Charig, A. J. & Milner, A. C. *Baryonyx*, a remarkable new theropod dinosaur. *Nature* **324**, 359–361 (1986).
- Schoener, T. W. The newest synthesis: understanding the interplay of evolutionary and ecological dynamics. *Science* **331**, 426–429 (2011).
- Houssaye, A. “Pachyostosis” in aquatic amniotes: a review. *Integr. Zool.* **4**, 325–340 (2009).
- Houssaye, A., Sander, M. P. & Klein, N. Adaptive patterns in aquatic amniote bone microanatomy—more complex than previously thought. *Integr. Comp. Biol.* **56**, 1349–1369 (2016).
- Quemeneur, S., De Buffrenil, V. & Laurin, M. Microanatomy of the amniote femur and inference of lifestyle in limbed vertebrates. *Biol. J. Linn. Soc.* **109**, 644–655 (2013).
- Canoville, A., de Buffrenil, V. & Laurin, M. Microanatomical diversity of amniote ribs: an exploratory quantitative study. *Biol. J. Linn. Soc.* **118**, 706–733 (2016).
- Amson, E., de Muizon, C., Laurin, M., Argot, C. & de Buffrenil, V. Gradual adaptation of bone structure to aquatic lifestyle in extinct sloths from Peru. *Proc. R. Soc. B* **281**, 20140192 (2014).
- Grafen, A. The phylogenetic regression. *Philos. Trans. R. Soc. B* **326**, 119–157 (1989).
- Liem, K. F. Adaptive significance of intra- and interspecific differences in the feeding repertoires of cichlid fishes. *Am. Zool.* **20**, 295–314 (1980).
- Turner, A. H., Pol, D., Clarke, J. A., Erickson, G. M. & Norell, M. A. A basal dromaeosaurid and size evolution preceding avian flight. *Science* **317**, 1378–1381 (2007).
- Voeten, D. F. et al. Wing bone geometry reveals active flight in *Archaeopteryx*. *Nat. Commun.* **9**, 1319 (2018).
- Houssaye, A., Martin, F., Boisserie, J. R. & Lihoreau, F. Paleocological inferences from long bone microanatomical specializations in Hippopotamoidea (Mammalia, Artiodactyla). *J. Mamm. Evol.* **28**, 847–870 (2021).
- Amson, E. & Bibi, F. Differing effects of size and lifestyle on bone structure in mammals. *BMC Biol.* **19**, 87 (2021).
- Malafafa, E. et al. A new spinosaurid theropod (Dinosauria: Megalosauroida) from the upper Barremian of Vallibona, Spain: Implications for spinosaurid diversity in the Early Cretaceous of the Iberian Peninsula. *Cret. Res.* **106**, 104221 (2020).
- Sereno, P. C. et al. A long-snouted predatory dinosaur from Africa and the evolution of spinosaurids. *Science* **282**, 1298–1302 (1998).
- Aureliano, T. et al. Semi-aquatic adaptations in a spinosaur from the Lower Cretaceous of Brazil. *Cret. Res.* **90**, 283–295 (2018).
- Barker, C. T. et al. New spinosaurids from the Wessex Formation (Early Cretaceous, UK) and the European origins of Spinosauridae. *Sci. Rep.* **11**, 19340 (2021).
- Taquet, P. *Géologie et Paléontologie du Gisement de Gadoufaoua (Aptien du Niger)* (Editions du Centre national de la Recherche Scientifique, 1976).
- Rayfield, E. J., Milner, A. C., Xuan, V. B. & Young, P. G. Functional morphology of spinosaur ‘crocodile-mimic’ dinosaurs. *J. Vertebr. Paleontol.* **27**, 892–901 (2007).
- Benson, R. B., Butler, R. J., Carrano, M. T. & O’Connor, P. M. Air-filled postcranial bones in the theropod dinosaurs: physiological implications and the ‘reptile–bird transition’. *Biol. Rev.* **87**, 168–193 (2012).

Publisher’s note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

© The Author(s), under exclusive licence to Springer Nature Limited 2022

Methods

Osteohistological analyses and assessment of skeletal maturity of spinosaurid specimens

Q13

We sampled and investigated postcranial elements of the holotype of *Baryonyx* (NHMR 9951), two individuals of *Suchomimus* (G51, G94), and the neotype of *Spinosaurus* (FSAC-KK 11888) to evaluate their somatic maturity and quantify bone density. The dorsal rib of *Baryonyx* was previously sectioned by Reid⁴⁴ and this was here studied for estimation of somatic maturity of the holotype. To quantify bone compactness, the femur of *Baryonyx* was computerized tomography (CT) scanned at the Natural History Museum, London. Breaks of additional postcranial bones (scapula, pubis, ischium and fibula) were considered for comparative purposes with the goal of assessing the skeletal extent of osteosclerosis. Skeletal maturity and bone compactness of *Suchomimus* were estimated through sampling and thin sectioning of the femora of the two individuals (described in brief in Ibrahim et al.⁹). A dorsal neural spine, dorsal rib, femur, and fibula were sampled for thin sectioning and inference of ontogenetic stage of the *Spinosaurus* neotype. Moreover, a manual phalanx, caudal neural spines and tibia of this specimen were also available for bone compactness quantification, because of breaks along the diaphysis. Long bones were cut transversely at the diaphysis, whereas samples of dorsal ribs and the dorsal neural spine of *Spinosaurus* were taken proximally and apically, respectively. The thin sectioning was performed following the protocol by Chinsamy & Rath⁴⁵. The thin sections have a thickness of 50–70 microns and were analysed with a Leica DM 2500 P petrographic microscope. Photographs of the bone tissue were taken with a ProgRes Cfsan camera. The CT scanned femur of *Baryonyx* was analysed in VGStudio Max 3.4. Inference of skeletal maturity follows recently proposed nomenclature by Griffin et al.⁴⁶.

Phylogenetic analyses

The discovery, description, and completeness of the *Spinosaurus* neotype provides an opportunity to revisit the phylogenetic relationships of spinosaurids. We coded the neotype of *Spinosaurus* in two recent datasets, published by Malafaia et al.³⁷ and Rauhut and Pol¹⁷, respectively. These two datasets differ in terms of operational taxonomic units (OTUs): whereas Malafaia et al.³⁷ includes a specimen-level assessment of phylogenetic relationships among spinosaurids, Rauhut and Pol¹⁷ remains the most comprehensive and latest iteration of the original dataset of tetanuran phylogenetic relationships³⁷. These two datasets are therefore needed to infer phylogenetic relationships within Spinosauridae and the placement of this clade among tetanurans. The neotype of *Spinosaurus* was coded as a separate OTU in the dataset of Malafaia et al.³⁷. On the basis of the results of this analysis, and given the presence of several apomorphies shared between the *Spinosaurus* neotype and holotype, coding of the neotype specimen based on our anatomical observations was added to the OTU of *Spinosaurus* in the dataset of Rauhut and Pol¹⁷. The recently described spinosaurine *Valibonavenatrix*³⁷ was also added to the dataset published by Rauhut and Pol¹⁷. We followed the most recent, comprehensive taxonomic and systematic revision of spinosaurid taxa^{48,49}, therefore excluding *Sigilmassasaurus* and *Oxalaia* (which are regarded as junior synonyms of *Spinosaurus*) from this dataset. Both datasets^{17,37} were analysed under equally weighted parsimony in TNT (Tree analysis using New Technology) v. 1.1⁵⁰. No characters were ordered. We conducted a heuristic search using 1,000 replicates of Wagner trees (with random addition sequence) followed by tree bisection and reconnection (TBR) branch swapping. We calculated decay indices (i.e., Bremer support) and absolute bootstrap frequencies with 10,000 pseudoreplicates to quantify node support.

Bone density

Bone density was used as a proxy for ecological inference. Because different postcranial skeletal elements show contrasting compactness

profiles due to allometry during growth^{15,27,51–54}, we focused on the femur and dorsal ribs in order to employ a consistent comparative framework; these skeletal elements have been previously shown to be reliable skeletal element for confident inference of ecological adaptations (for example, in ref.²⁷). Femoral and dorsal rib cross sections were mainly obtained from the diaphysis and the proximal region, respectively, through thin sectioning, micro-CT scanning, or data mining from the literature (see Supplementary Dataset for the taxa included and the type of data collected). Among dinosaurs, novel data presented in this study include those for the tetanuran theropods *Baryonyx*, *Suchomimus*, *Spinosaurus*, *Megalosaurus*, *Tyrannotitan*, *Eustreptospondylus*, and *Condorraptor* (see Supplementary Dataset for novel osteohistological data collected for this study). Our femoral dataset includes 206 individuals, representing 200 taxa. All known spinosaurid taxa that preserve the femur are included therein. The discrepancy between the number of individuals and taxa is due to the inclusion of multiple individuals of the following marine reptiles: *Ichthyosaurus*, *Nothosaurus*, *Simosaurus*, Placodontia and *Champsosaurus*. Our dorsal rib dataset includes 174 taxa. The taxonomic overlap between the two datasets (femur and dorsal rib) is equal to 83 taxa, including *Baryonyx* and *Spinosaurus*.

Archosaurs are represented in the dataset by extant crocodylians, pterosaurs, non-avian dinosaurs, and birds, the latter including both Mesozoic and extant taxa (see Supplementary Dataset for included taxa). Stem and crown marine mammals, such as cetaceans and seals, and extinct marine reptiles (ichthyosaurs, sauropterygians, and mosasaurs) were included to infer thresholds of bone compactness related to aquatic lifestyle and to calibrate the discriminant analyses aimed to infer ecological adaptations in extinct taxa.

Cross (CT scan) and thin sections of femoral diaphysis and dorsal rib were transformed into black and white figures (black for bone and white for medullary cavity, vascularization, and background) in Adobe Photoshop, following previous protocols (for example, refs.^{15,27,36,54}). Images were then imported into the freely available software Bone Profiler⁵⁵ (<http://134.158.74.46/BoneProfileR/>) to quantify bone compactness. In cases where portions of the femoral diaphysis and rib cross sections were missing or deformed, retro-deformation and reconstruction were applied following the methods presented by De Ricqlès et al.⁵⁶, to minimize the occurrence of taphonomic artifacts in the data. Because the femoral diaphysis of *Baryonyx* is eroded and crushed, the cross section for this taxon was taken from a more intact and better-preserved region closer to the distal portion of the femur (Supplementary Fig. 3). Because the diaphysis of the femur coincides with the highest degree of bone compactness among amniotes^{15,27}, the quantified degree of osteosclerosis in *Baryonyx* should be regarded as underestimated.

Informal consensus tree

To address the statistical non-independence of interspecific comparisons, we assembled two informal amniote-wide supertrees (Extended Data Figs. 8, 9) using Mesquite v. 3.40⁵⁷ on the basis of Upham et al.⁵⁸ for Mammalia, Simoes et al.⁵⁹ for the backbone of Diapsida, Nesbitt et al.⁶⁰ for Archosauria, Langer et al.⁶¹ for Dinosauria, this study for Tetanurae, Brusatte et al.⁶² for Coelurosauria, and Prum et al.⁶³ for Neoaves. We calibrated the resulting tree using the function 'bin_timePaleoPhy' from the R package Paleotree⁶⁴, scaling the branches on the basis of genus-level stratigraphic ranges sourced from the Paleobiology Database (www.paleodb.org) and from the specialized literature (see Supplementary Dataset). We generated 100 trees using this method, which randomly draws first appearance dates and last appearance dates for each taxon from within their stratigraphic ranges. To avoid zero-length branches we set a minimum branch length of one million years.

Ecological inference

We scored extant and extinct taxa whose ecomorphological attributes could confidently be inferred (for example, ichthyosaurs as being able

to dive frequently and not being able to fly) as being able to engage in (a) subaqueous-foraging (0, unable; 1, able but infrequent (for example, rails); 2, frequent), and (b) flying (0, unable; 1, non-sustained flight (for example, tinamous, galliforms and *Xenicus longipes*); 2, sustained flight). Extinct taxa with ambiguous ecological inference were scored as unknown. Therefore, the autecology of each taxon is represented by two numerical categories with three states each. Previous studies applied different categorizations for the characterization of aquatic lifestyles among extant and extinct taxa: ‘aquatic’ and ‘semiaquatic’ were used as the opposite of “subaqueous foraging” applied in this study. Our ecomorphological attribution is focused on a specific behaviour linked to an ecology, rather than a categorization of its entirety. We find our categorization to be more accurate: for example, previous studies coded penguins and cetaceans as aquatic, while crocodylians were stated as semiaquatic. Whereas penguins and crocodylians are still ecologically dependent on terrestrial environments (for example, for laying eggs), cetaceans are completely independent from land. On the other hand, all these clades engage in subaqueous foraging. Therefore, our ecological attribution is in agreement with previously applied ecological categories, but do not exclude dependency to terrestrial environments to satisfy autecological requirements, such as reproductive behaviour.

Maximum femoral diaphyseal and dorsal rib cross section diameter was used as a proxy for body size, in order to allow the inclusion of fragmentary fossil remains and to optimize the inclusion of taxa with significantly different body plans. As femoral and rib diameter values range from those of small-bodied modern passerines (*Xenicus*) to very large non-avian theropods (*Tyrannosaurus* and *Spinosaurus*), maximum femoral diameter was log₁₀-transformed.

Bone compactness, femoral midshaft diameter, and different combinations of these ecological traits were used to build 12 linear models upon which PGLS were performed using the R core function `gls` (R Core Team). The AIC was used to establish which linear model best explains variation in bone compactness. Pagel’s lambda values were simultaneously calculated to evaluate the degree of phylogenetic signal in each of the relationships. These analyses were run over the 100 trees generated for all amniotes to evaluate the effects of stratigraphic uncertainty on our analyses; the results were summarized thereafter.

To establish explicit predictions of ecology in extinct taxa, we built a phylogenetically flexible discriminant analysis (pfDA) using the function `phylo.fda` (Schmitz & Motani⁶⁵, sourced from <https://github.com/lSchmitz/phylo.fda>) and following the protocol described⁶⁵, including our two main metric variables (maximum diameter and bone density) and the ecological classifiers from the linear model with the best fit (lowest AIC score). The model in which bone density is explained by subaqueous-foraging exhibits the best fit (see Results, Table 1, and Supplementary Tables 3, 4); therefore, we scored all taxa to a more inclusive category depending on whether or not they are frequent subaqueous-foragers (Supplementary Tables 3, 4). Because our overarching goal is to ascertain aquatic proficiency in large, flightless theropod dinosaurs, we also excluded modern birds that are able to both submerge-forage and fly as this functional trade-off is likely to influence their bone histology and introduce a confounding factor in our predictions. A series of taxa for which aquatic lifestyles have been proposed or fragmentary remains cannot allow a confident scoring were scored as ‘unknown’ and their ecologies were predicted along with the three spinosaurid target taxa (see supplementary dataset). In order to correct for the bias that phylogenetic structure introduces in form to function relationships, `phylo.fda` adjusts the phylogeny with the value of phylogenetic signal (Pagel’s lambda) which maximizes the log likelihood of the linear fit among variables^{65,66}. Because branch lengths in our phylogenies exhibit some degree of uncertainty, we repeated this analysis with the 100 different trees we generated and summarized the accuracy and predictions across all iterations. This was repeated for both the femora and rib datasets and again excluding graviportals and deep diving taxa in both datasets (see Supplementary Tables 5, 6

for taxa classified with these ecological traits). In each iteration, the variables (bone compactness and diameter) from the training set of taxa with known ecologies, together with the phylogenetic structure of data, are used to generate the discriminant functions, which are subsequently used to predict the ecologies in extinct taxa with unknown ecologies (including spinosaurids). A said species is predicted as subaqueous forager if the posterior probability is 50% or more, because our inference has only two possible outcomes: subaqueous forager or non-subaqueous forager. We summarised our results by providing the median value of those 100 posterior probabilities and the number of times a particular taxon is predicted as subaqueous forager (median probability of 50% or more). This gives us two proxies of the likelihood of each taxon to be an actual subaqueous forager. For instance, a taxon could be predicted 100 times as subaqueous forager with a median probability of 51% which means the evidence for this extinct species to be an actual subaqueous forager is very weak and this inference has to be considered very unlikely. Median probabilities need to be within the range of 80–100% to be considered as strong evidence of subaqueous forager. Additionally, we considered the presence of an open medullary cavity or osteosclerosis to support our inferences.

Reporting summary

Further information on research design is available in the Nature Research Reporting Summary linked to this paper.

Data availability

All data described and used in this manuscript are freely available. The measurements and provenance information for fossil specimens can be found in the extended data figures and in the Supplementary Dataset. The phylogenetic datasets and the R coding are available as Supplementary Material. The CT scan datasets collected for this study are available in Morphosource (specific links for each taxon can be found in the Supplementary Dataset).

44. Reid, R. E. H. Zonal “growth rings” in dinosaurs. *Mod. Geol.* **15**, 19–48 (1990).
45. Chinsamy, A. & Raath, M. A. Preparation of fossil bone for histological examination. *Palaeont. Afr.* **29**, 39–44 (1992).
46. Griffin, C. T. et al. Assessing ontogenetic maturity in extinct saurian reptiles. *Biol. Rev.* **96**, 470–525 (2021).
47. Carrano, M. T., Benson, R. B. & Sampson, S. D. The phylogeny of Tetanurae (Dinosauria: Theropoda). *J. Syst. Palaeontol.* **10**, 211–300 (2012).
48. Ibrahim, N. et al. Geology and paleontology of the Upper Cretaceous Kem Kem Group of eastern Morocco. *ZooKeys* **928**, 1–216 (2020).
49. Smyth, R. S., Ibrahim, N. & Martill, D. M. *Sigilmassasaurus* is *Spinosaurus*: a reappraisal of African spinosaurines. *Cret. Res.* **114**, 104520 (2020).
50. Goloboff, P. A., Farris, J. S. & Nixon, K. C. TNT, a free program for phylogenetic analysis. *Cladistics* **24**, 774–786 (2008).
51. Erickson, G. M. Assessing dinosaur growth patterns: a microscopic revolution. *Trends Ecol. Evol.* **20**, 677–684 (2005).
52. Hayashi, S. et al. Bone inner structure suggests increasing aquatic adaptations in Desmostylia (Mammalia, Afrotheria). *PLoS ONE* **8**, e59146 (2013).
53. Straehl, F. R., Scheyer, T. M., Forasiepi, A. M., MacPhee, R. D. E. & Sánchez-Villagra, M. R. Evolutionary patterns of bone histology and bone compactness in xenarthran mammal long bones. *PLoS ONE* **8**, e69275 (2013).
54. Houssaye, A., Tafforeau, P., de Muizon, C. & Gingerich, P. D. Transition of Eocene whales from land to sea: evidence from bone microstructure. *PLoS ONE* **10**, e0118409 (2015).
55. Girondot, M. & Laurin, M. Bone profiler: a tool to quantify, model, and statistically compare bone-section compactness profiles. *J. Vertebr. Paleontol.* **23**, 458–461 (2003).
56. De Ricqlès, A. J., Padian, K., Horner, J. R., Lamm, E. T. & Myhrvold, N. Osteohistology of *Confuciusornis sanctus* (Theropoda: Aves). *Journ. Vertebr. Paleontol.* **23**, 373–386 (2003).
57. Maddison, W. P. Mesquite: a modular system for evolutionary analysis. *Evolution* **62**, 1103–1118 (2008).
58. Upham, N. S., Esselstyn, J. A. & Jetz, W. Inferring the mammal tree: species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS Bio.* **17**, e3000494 (2019).
59. Simoes, T. R. et al. The origin of squamates revealed by a Middle Triassic lizard from the Italian Alps. *Nature* **557**, 706–709 (2018).
60. Nesbitt, S. J. et al. The earliest bird-line archosaurs and the assembly of the dinosaur body plan. *Nature* **544**, 484–487 (2017).
61. Langer, M. C. et al. Untangling the dinosaur family tree. *Nature* **551**, E1–E3 (2017).
62. Brusatte, S. L., Lloyd, G. T., Wang, S. C. & Norell, M. A. Gradual assembly of avian body plan culminated in rapid rates of evolution across the dinosaur-bird transition. *Curr. Biol.* **24**, 2386–2392 (2014).

63. Prum, R. O. et al. A comprehensive phylogeny of birds (Aves) using targeted next-generation DNA sequencing. *Nature* **526**, 569–573 (2015).
64. Bapst, D. W. paleotree: an R package for paleontological and phylogenetic analyses of evolution. *Methods Eco. Evo.* **3**, 803–807 (2012).
65. Schmitz, L. & Motani, R. Nocturnality in dinosaurs inferred from scleral ring and orbit morphology. *Science* **332**, 705–708 (2011).
66. Motani, R. & Schmitz, L. Phylogenetic versus functional signals in the evolution of form–function relationships in terrestrial vision. *Evolution* **65**, 2245–2257 (2011).

Acknowledgements We acknowledge P. Barrett and S. Chapman for access to the holotype of *Baryonyx* at the Natural History Museum, London, J. Scannella for access to thin sections of *Tyrannosaurus* housed at the Museum of the Rockies, and M. Fox and J. Gauthier for access to *Poposaurus* at the Yale Peabody Museum. The Moroccan Ministry of Energy, Mines, and the Environment is thanked for providing fieldwork permits to N. I. Members of the 2015–2019 expedition seasons are thanked for their assistance in the field. We thank J. Choiniere, three anonymous reviewers, and the editor for constructive comments that improved the manuscript. Funding was received from the European Union's Horizon 2020 research and innovation program 2014–2018, starting grant (R. B. J. B., 677774); a National Geographic Society grant (N.I., CP-143R-170); a National Geographic Emerging Explorer Grant (N.I.); the Jurassic Foundation (M.F.); the Paleontological Society grant (M.F.), as well as the Explorers

Club (grant awarded to M.F.). The Lokschuppen (Rosenheim, Germany) and J. Pfauntsch provided additional financial support for fieldwork led by N.I. in Morocco.

Author contributions Conceptualization: M.F. Data collection and curation: all authors. Data quantification: M.F. Methodology: M.F., G.N. and R.B.J.B. Formal analysis: M.F., G.N. and R.B.J.B. Resources: all authors. Writing, original draft preparation: M.F. Writing, review and editing: all authors. Visualization: M.F. and G.N.; Supervision: M.F., G.N., R.B.J.B. and N.I. Funding acquisition: M.F., R.B.J.B. and N.I.

Competing interests The authors declare no competing interests.


Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41586-022-04528-0>.

Correspondence and requests for materials should be addressed to Matteo Fabbri, Guillermo Navalón or Roger B. J. Benson.


Peer review information *Nature* thanks Jonah Choiniere, Peter Falkingham, Sterling Nesbitt and the other, anonymous, reviewers for their contribution to the peer review of this work. Peer review reports are available.

Reprints and permissions information is available at <http://www.nature.com/reprints>.




Extended data Fig. 1 | Comparative array of archosaurian femoral diaphysis included in the dataset. Numerical values represent the bone density quantified for each taxon. Asterisks indicate femoral diaphysis that were


retro-deformed before quantification of bone density due to taphonomic deformation and/or fragmentation present in the fossil.



Extended data Fig. 2 | Comparative array of non-avian and avian femoral diaphysis included in the dataset. Numerical values represent the bone density quantified for each taxon.




Extended data Fig. 3 | Comparative array of avian and lepidosaur femoral diaphysis included in the dataset. Numerical values represent the bone density quantified for each taxon.




Extended data Fig. 4 | Comparative array of amniote femoral diaphysis included in the dataset. Numerical values represent the bone density quantified for each taxon.

Article




Extended data Fig. 5 | Comparative array of mammalian femoral diaphysis included in the dataset. Numerical values represent the bone density quantified for each taxon.



Extended data Fig. 6 | Comparative array of archosaurian dorsal rib cross sections included in the dataset. Numerical values represent the bone density quantified for each taxon.


Article



Extended data Fig. 7 | Comparative array of amniote dorsal rib cross sections included in the dataset. Numerical values represent the bone density quantified for each taxon.


Femur midshaft diameter (log10 mm)

-0.036 2.672



Bone compactness

0.279 0.989




Extended Data Fig. 8 | Bone density and femur diameter phylogenetic distribution plotted on the informal consensus tree used for discriminant analyses representing the phylogenetic relationships of the taxa included in our study.

Ribs midshaft diameter (log10 mm)

Bone compactness

-0.378 1.978

545.0 866.0




Extended Data Fig. 9 | Bone density and dorsal rib diameter phylogenetic distribution plotted on the informal consensus tree used for discriminant analyses representing the phylogenetic relationships of the taxa included in our study.

Scapula

Baryonyx

bd=0.954



Rahonavis


bd=0.602



Dorsal rib


Baryonyx

bd=0.921



Saltriovenator

bd=0.704



Dorsal neural spine


Spinosaurus

bd=0.929



Ouranosaurus

bc=0.58



Tibia


Spinosaurus

bd=0.946



Tyrannosaurus

bd=0.692



Extended data Fig. 10 | Qualitative comparison of bone compactness in selected skeletal elements between osteosclerotic spinosaurids and other non-avian dinosaurs. *Baryonyx* and *Spinosaurus* possess dense, compact bone throughout the postcranial skeleton, namely in the neural spines, ribs,

scapula, ilium, pubis, ischium, femur, and fibula. Increased bone density is found in postcranial elements of *Spinosaurus* as well; a reduced medullary cavity is present in the ribs, dorsal and caudal neural spines, manual phalanges, femur, tibia, and fibula. Abbreviations: bd=bone density.

Author Queries

Journal: **Nature**

Paper: **s41586-022-04528-0**

Title: **Subaqueous foraging among carnivorous dinosaurs**

AUTHOR:

The following queries have arisen during the editing of your manuscript. Please answer by making the requisite corrections directly in the e-proofing tool rather than marking them up on the PDF. This will ensure that your corrections are incorporated accurately and that your paper is published as quickly as possible.

Query Reference	Reference
Q1	Please provide Websum.
Q2	Your paper has been copy edited. (1) Please review every sentence to ensure that it conveys your intended meaning; if changes are required, please provide further clarification rather than reverting to the original text. Please note that formatting (including hyphenation, Latin words, and any reference citations that might be mistaken for exponents) and usage have been made consistent with our house style. (2) Check the title and the first paragraph with care, as they may have been re-written to aid accessibility for non-specialist readers. (3) Check the symbols for affiliations with care, and check all author names and Acknowledgements carefully to ensure that they are correct; check the email address of the corresponding author and the Competing Interests statement. (4) Check that there has been no corruption of mathematical symbols. Single-letter variables are set in italics (but not their subscripts unless these are also variables). Vectors are set as bold; matrices are set as italic only. We do not use italics for emphasis. Genetic material is set in italic and gene products are set upright. Please check that italicization and bolding are correct throughout. (5) Ensure that, where practicable, all figures, tables and other discrete elements of Supplementary Information are referred to at least once in the paper at an appropriate place in the text or figure legends. (6) Please note, we reserve 'significant' and its derivatives for statistical significance. Please reword where this is not the intended meaning (for example, to important, notable, substantial).
Q3	Please note that the eproof should be amended in only one browser window at any one time, otherwise changes will be overwritten.
Q4	Please check that the display items are as follows (ms no: 2021-07-11463C): Figs none (black & white); 3 (colour); Tables: 1; Boxes: None; Extended Data display items: 10; SI: yes. The eproof contains the main-text figures edited by us and (if present) the Extended Data items (unedited except for minor formatting) and the Supplementary Information (unedited). Please check the edits to all main-text figures (and tables, if any) very carefully, and ensure that any error bars in the figures are defined in the figure legends. Extended Data items may be revised only if there are errors in the original submissions. If you need to revise any Extended Data items please upload these files when you submit your corrections to this eproof, and include a list of what has been changed.

Author Queries

Journal: **Nature**

Paper: **s41586-022-04528-0**

Title: **Subaqueous foraging among carnivorous dinosaurs**

AUTHOR:

The following queries have arisen during the editing of your manuscript. Please answer by making the requisite corrections directly in the e-proofing tool rather than marking them up on the PDF. This will ensure that your corrections are incorporated accurately and that your paper is published as quickly as possible.

Query Reference	Reference
Q5	A single sentence summarizing your paper has been provided (editor's summary in the eproof), which will appear online on the table of contents and in e-alerts. Please check this sentence for accuracy and appropriate emphasis.
Q6	Please check your article carefully, coordinate with any co-authors and enter all final edits clearly in the eproof, remembering to save frequently. Once corrections are submitted, we cannot routinely make further changes to the article.
Q7	Note that the eproof should be amended in only one browser window at any one time; otherwise changes will be overwritten.
Q8	Author surnames have been highlighted. Please check these carefully and adjust if the first name or surname is marked up incorrectly. Note that changes here will affect indexing of your article in public repositories such as PubMed. Also, carefully check the spelling and numbering of all author names and affiliations, and the corresponding email address(es).
Q9	You cannot alter accepted Supplementary Information files except for critical changes to scientific content. If you do resupply any files, please also provide a brief (but complete) list of changes. If these are not considered scientific changes, any altered Supplementary files will not be used, only the originally accepted version will be published.
Q10	If applicable, please ensure that any accession codes and datasets whose DOIs or other identifiers are mentioned in the paper are scheduled for public release as soon as possible, we recommend within a few days of submitting your proof, and update the database record with publication details from this article once available.

Author Queries

Journal: **Nature**

Paper: **s41586-022-04528-0**

Title: **Subaqueous foraging among carnivorous dinosaurs**

AUTHOR:

The following queries have arisen during the editing of your manuscript. Please answer by making the requisite corrections directly in the e-proofing tool rather than marking them up on the PDF. This will ensure that your corrections are incorporated accurately and that your paper is published as quickly as possible.

Query Reference	Reference
Q11	<p>(1) Please ensure that the following information is included in the figure legends where relevant. Sample size (exact n number); a statement of replicability (how many times was experiment replicated in the lab); description of sample collection (clarify whether technical or biological replicates and include how many animals, litters, cultures, etc.); state the statistical test used and give P values; define centre values (median or average) and error bars. (2) For figures/images that are reproduced or adapted from a third party, it is important that you confirm that permission has been obtained and that appropriate acknowledgement of the copyright holder is given. (3) Please note that we edit the main figures (but not the Extended Data figures) in house. There is no need to resupply any of the main figures to make minor changes to text labels to match the changes made in the text, as figures will have been edited accordingly. If you wish to make changes to any Extended Data figures, however, please resupply these, and please let us know what has changed.</p>
Q12	<p>Fig. 2a and 3a legends describe a dashed line, but there is no dashed line in the figure. Please check.</p>
Q13	<p>(1) Please ensure that the following information is provided in the Methods section where relevant. Animal experiments require: statement about randomization; statement about blinding; statement of sex, age, species and strain of animals; statement of IRB approval for live vertebrate experimentation. For experiments involving humans: statement of IRB approval; statement of informed consent; statement of consent to publish any photos included in figures. Randomized clinical trials require trial registration. (2) We recommend that detailed protocols are deposited in Protocol Exchange, or a similar repository. (3) If custom computer code has been used and is central to the conclusions of this paper, please insert a section into the Methods titled 'Code availability' and indicate within this section whether and how the code can be accessed, including any restrictions to access. (4) Please state whether statistical methods were used to predetermine sample size. (5) Please state whether blinding and randomization were used. (6) To address the issue of cell line misidentification and cross-contamination, for any cell lines mentioned in the paper please provide source of the cell lines and indicate whether the cell lines have been correctly identified/authenticated (if so, by what methods). Also, please state whether cell lines have been tested for mycoplasma contamination.</p>

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- A description of all covariates tested
- A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection

Femoral diaphysis and dorsal rib cross sections belonging to 206 and 174 amniote, respectively, were collected through means of microCT scan or thin sectioning (e.g. Spinosaurus, Suchomimus, and Baryonyx), or obtained from the literature. Sections were then imaged to allow quantification of bone compactness and quantitative analyses. See the supplementary data attached to this manuscript for a complete list of data used in this study, origin of the data, and type of data acquisition. Femoral and dorsal rib sections were converted into black and white figures in VGStudio Max 3.4 for data obtained through microCT scan or in Photoshop CS5 for imaged thin sections. Bone density was quantified with the freely available software BoneProfiler 2.0-1 (<https://max2.ese.u-psud.fr/epc/conservation/boneprofiler.html>). The freely available software "Fiji" (<https://imagej.net/software/fiji/>) was used to measure the maximum diameter of the femur diaphyses and rib cross sections.

Data analysis

Phylogenetic analyses were performed in TNT (Tree analysis using New Technology) v. 1.1. In order to build an informal Supertree to test the role of phylogeny on the evolution of bone density, we used the freely available software Mesquite 3.7 (<https://www.mesquiteproject.org/>). The tree was calibrated using "bin_timePaleoPhy" from the R package Paleotree. Bone density, femoral midshaft and rib cross section diameters, and different combinations of ecological traits were used to build 12 linear models upon which phylogenetically gnostic regressions (PGLS) were performed using the R core function gls (R Core Team). To establish explicit predictions of ecology in extinct taxa, we built a phylogenetically-flexible discriminant analysis (pfDA) using the function phylo.fda (sourced from <https://github.com/lSchmitz/phylo.fda>). In order to correct for the bias that phylogenetic structure introduces in form to function relationships, phylo.fda adjusts the phylogeny with the value of phylogenetic signal (Pagel's lambda) which maximizes the log likelihood of the linear fit among variables.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

All data described in this paper, including measurements, provenance information for fossil specimens, phylogenetic datasets, and CT scans collected for this study, are available in the Extended Data Figures, Supplementary Materials, and in Morphosource

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

- Life sciences Behavioural & social sciences Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	In this study, we conduct phylogenetic comparative analyses of osteohistological and bone density data in a broad sample of amniotes, including dinosaurs, (n=380) to assess the presence and extent of aquatic adaptations in non-avian dinosaurs. First, we use comparative analysis to validate bone compactness as a proxy for subaqueous foraging capabilities in amniotes. We then use this proxy to provide evidence that a clade of medium- to giant-bodied predatory dinosaurs, Spinosauridae, was ecologically adapted to life in water, representing the first known aquatic radiation among non-avian dinosaurs. Disparate ecological niches are herein recognized among spinosaurids: whereas Baryonyx and Spinosaurus are strongly predicted as subaqueous foraging species, Suchomimus is inferred as a more terrestrial, potentially wading animal, although sharing morphological similarities with Baryonyx. Our findings greatly expand the ecological disparity of non-avian dinosaurs.
Research sample	This study collected bone compactness values from the femora (n=206) and dorsal ribs (n=174) of extant and extinct amniote taxa. The data collected aimed to cover the broadest taxonomic and ecological diversity possible among amniotes. While part of the data were available from the literature, our team collected data for 79 modern and extinct taxa.
Sampling strategy	We sampled 374 amniote taxa (380 individuals) spanning the entire tree of life, body size variation, and ecological diversity of this clade, in order to verify the influence of phylogeny, allometry and ecology on bone density, and to infer ecological adaptations among non-avian dinosaurs. The dataset here published is the largest ever built for bone density in amniotes. Because this includes extant and extinct clades and captures a broad ecological variability within each major amniote clade, we believe that the analyzed dataset is a consistent representation of ecological evolution in Amniota.
Data collection	Data collected for this study were acquired through thin sectioning or microCT scan. These were collected in the respective Institutions housing the specimens. MicroCT scans were collected for the diaphyses of femora and then imported in VGStudio Max 3.4 where a section of the diaphysis was extracted. Thin sections were imaged through petrographic microscopes and then transformed in Photoshop in a black and white image (black for bone). See the supplementary dataset attached to this study for the list of taxa examined. In order to expand the dataset and cover a broader range of taxa and ecologies, we obtained data from the literature. In this case, thin sections were directly transformed in black and white figures in Photoshop. In few cases, quantification of bone density was already undertaken in literature: in these cases, the values for bone density and maximum diameter were simply collected. While all authors contributed with specimens, M.F. transformed the sections in black and white figures and quantified bone density and maximum diameter of the sections.
Timing and spatial scale	Data were collected from FMNH, YPM at Yale, AMNH, NHMUK, UMZC, SMNS, UUVF, IGWH, NMQR, NMW, BSP, MPEF, AODF, IVPP, UM, MBC, MOR, OUMNH, MWU, MLP, STIPB, CAV, GPIT, UMMZ, OUMNH-J, LACM, SMNS, IPS, LO, RR, TMM, ROM, Wijk, SIPB R, ZMNH, CRILAR, UA, IPB, GAD, FSAC-KK, BPI, UF, ML, and MACN-PV. Data collection started in 2015 and was constantly conducted until 2019.
Data exclusions	No data were excluded
Reproducibility	Quantification of bone density was performed three times for each taxon to confidently replicate density measurements. Analyses of ecological inference were run over 100 trees generated for all amniotes to evaluate the effects of stratigraphic uncertainty on our analyses; the results were summarized thereafter. All data and coding required to replicate our results are available as supplementary material to this study. Replication of the results were successful.
Randomization	Randomization was applied to the alternative 100 phylogenies obtained for the PGLS analyses aimed at finding potential correlations between bone density, ecology and allometry. All other analyses do not require randomization.
Blinding	Not applicable because experiments do not require blinding

Did the study involve field work? Yes No

Field work, collection and transport

Field conditions	Desert escarpment close to Moroccan-Algerian border
Location	In- and ex-situ on the slopes of a south-east facing escarpment (Zrigat, southeastern Morocco; 31° 37' N, 4° 16' W) fringing the Aferdou Zrigat plateau (Tafilalt basin, Akrabou Formation, Kem Kem beds).
Access & import/export	Permits for fieldwork were obtained from Ministère de l'Energie, des Mines, et de l'Environnement. Permits: 4581/DE/2019 (issued on 17/07/2019) and 4118/DE/2018/DG (issued on 06.06.2018). The work was performed in close collaboration with researchers in Morocco (FSAC, Casablanca). The specimens collected are deposited at the Departement de Géologie/Laboratoire de Biodiversité et Santé, Faculté des Sciences Ain Chock, Hassan II University, Casablanca, Morocco.
Disturbance	No disturbance

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> Antibodies
<input checked="" type="checkbox"/>	<input type="checkbox"/> Eukaryotic cell lines
<input type="checkbox"/>	<input checked="" type="checkbox"/> Palaeontology and archaeology
<input checked="" type="checkbox"/>	<input type="checkbox"/> Animals and other organisms
<input checked="" type="checkbox"/>	<input type="checkbox"/> Human research participants
<input checked="" type="checkbox"/>	<input type="checkbox"/> Clinical data
<input checked="" type="checkbox"/>	<input type="checkbox"/> Dual use research of concern

Methods

n/a	Involvement in the study
<input checked="" type="checkbox"/>	<input type="checkbox"/> ChIP-seq
<input checked="" type="checkbox"/>	<input type="checkbox"/> Flow cytometry
<input checked="" type="checkbox"/>	<input type="checkbox"/> MRI-based neuroimaging

Palaeontology and Archaeology

Specimen provenance	The 380 specimens used in the study range from Europe, Mongolia, Africa, North and South America.
Specimen deposition	The specimens used in this study are housed in public museum collections and are freely accessible. The Institutions where the investigated specimens are housed are listed in the supplementary data attached to this manuscript. Data were collected from FMNH, YPM at Yale, AMNH, NHMUK, UMZC, SMNS, UUV, IGWH, NMQR, NMW, BSP, MPEF, AODF, IVPP, UM, MBC, MOR, OUMNH, MWU, MLP, STIPB, CAV, GPIT, UMMZ, OUMNH-J, LACM, SMNS, IPS, LO, RR, TMM, ROM, Wijk, SIPB R, ZMNH, CRILAR, UA, IPB, GAD, FSAC-KK, BPI, UF, ML, and MACN-PV.
Dating methods	No new dates were provided
<input checked="" type="checkbox"/>	Tick this box to confirm that the raw and calibrated dates are available in the paper or in Supplementary Information.
Ethics oversight	No ethical oversight was needed for this study, because based on Museum specimens

Note that full information on the approval of the study protocol must also be provided in the manuscript.