



MINI REVIEW

Microbiome changes in esophageal cancer: implications for pathogenesis and prognosis

Yi Li^{1*}, Bing Wei^{1,2*}, Xia Xue³, Hongle Li^{1,2}, Jun Li^{1,2}

¹Department of Molecular Pathology, Clinical Pathology Center, Affiliated Cancer Hospital of Zhengzhou University and Henan Cancer Hospital, Zhengzhou 450003, China; ²Henan Key Laboratory of Molecular Pathology, Zhengzhou 450003, China; ³Henan Key Laboratory of Helicobacter pylori & Microbiota and Gastrointestinal Cancer, Marshall Medical Research Center, The Fifth Affiliated Hospital of Zhengzhou University, Zhengzhou 450052, China

ABSTRACT

Esophageal cancer (EC) is an aggressive malignancy with a poor prognosis. Various factors, including dietary habits, and antacid and antibiotic use, have been shown to influence the esophageal microbiome. Conversely, enrichment and diversity of the esophageal microbiome can also impact its function. Recent studies have revealed prevalent changes in the esophageal microbiome among patients with EC, thus suggesting the potential contribution of the esophageal microbiome to EC development. Additionally, distinct microbiome compositions have been observed in patients with different responses to radiotherapy and chemotherapy, indicating the role of the esophageal microbiome in modulating treatment outcomes. In this review, we have examined previous studies on the esophageal microbiome in healthy individuals and patients with EC or other esophageal diseases, with a focus on identifying microbial communities associated with EC pathogenesis and prognosis. Understanding the role of the microbiome in EC may aid in early detection and optimized treatment strategies, ultimately leading to better outcomes for patients.

KEYWORDS

Esophageal cancer; microbiome; dysbiosis; microenvironment; carcinogenesis

Introduction

Esophageal cancer (EC) is the seventh most common cancer and the sixth leading cause of cancer-related deaths worldwide¹. Esophageal squamous cell carcinoma (ESCC) and esophageal adenocarcinoma (EAC) are the two primary subtypes of EC, but ESCC and EAC differ significantly with respect to geographic patterns, temporal trends, and risk factors^{2,3}. ESCC is prevalent in central Asia and the easterly-lying corridor in Africa, extending from the Taihang Mountains in northern China to central Asia, Iran, and eastern and southern Africa². In contrast, EAC is more prevalent in industrialized countries in Europe³.

The symptoms of EC are correlated with disease progression. In the early stages, patients may be asymptomatic or complain of dysphagia, while in the advanced stages patients may exhibit progressive dysphagia, persistent retrosternal or back pain, and marked cachexia. Endoscopic therapy is routinely used to remove cancerous tissues in early-stage disease. Chemotherapy and radiation therapy are usually applied before or after surgical resection for locally advanced disease⁴, while treatment options for metastatic disease include chemotherapy, radiation therapy, targeted therapy, immunotherapy, or a combination of these treatments to manage symptoms and slow disease progression. The use of neoadjuvant radiotherapy and chemotherapy during the perioperative period can effectively prolong the median survival rate for patients with advanced disease, particularly patients with ESCC. Surgical resection after neoadjuvant radiotherapy and chemotherapy has become the standard treatment for locally advanced ESCC⁵.

Several factors are associated with an increased risk of EC (Table 1). The incidence of ESCC is typically higher in men, while women may be more susceptible to developing EAC; however, the exact mechanisms underlying this difference

*These authors contributed equally to this work.

Correspondence to: Hongle Li and Jun Li

E-mail: llhl73@163.com and zlylijun3922@zzu.edu.cn

ORCID ID: <https://orcid.org/0000-0001-6428-9091>

and <https://orcid.org/0000-0001-8815-3578>

Received May 19, 2023; accepted September 6, 2023;

published online October 9, 2023.

Available at www.cancerbiomed.org

©2023 Cancer Biology & Medicine. Creative Commons

Attribution-NonCommercial 4.0 International License

Table 1 Known risk factors for esophageal carcinoma

Exposure	Outcomes
Gender	For EAC and ESCC, males have a higher risk than females ⁶
Tobacco	Smoking cigarettes, pipes, cigars, hookah, and chewing tobacco are high-risk factors for EC ⁷
Alcohol	Alcohol is not a risk factor for EAC and is not closely related to ESCC ⁸
Diet	Pickled vegetables and hot foods are historically related to the high incidence of EC ⁹
Fat	Central obesity is a major and consistent risk factor for the development of EAC ¹⁰
Social status	Poverty and a low level of education are among the most dangerous risk factors for EC ¹¹

are unclear⁶. Smokers have a five-fold higher risk of EC than non-smokers⁷. Alcohol consumption does not appear to be a significant risk factor for both ESCC and EAC². Obesity, particularly central obesity, may increase intra-abdominal and gastric pressure, promote gastroesophageal reflux, and promote the development of EAC¹⁰. Individuals residing in developing countries or regions with limited access to fruits and vegetables have increased susceptibility to esophageal inflammation and cancer². Individuals with low levels of education, physical activity, and income have a higher risk of EAC in developed countries; currently, a similar trend has been noted in developing countries^{9,11}. More recently, researchers have studied alterations in esophageal microbiota that may contribute to the development of EC.

Culture-independent high-throughput DNA sequencing technologies, coupled with advanced computational tools, empower the comprehensive analysis of the intricate human microbiome¹¹. It is well-established that the gut microbiome, which consists of trillions of microorganisms, exerts a significant influence on facilitating the development of gastrointestinal tumors and suppressing antitumor immune responses¹²; however, the potential contribution of esophageal microorganisms to the development of EC has yet to be fully elucidated. One of the hurdles in studying the esophageal microbiome involves obtaining samples from the esophageal epithelium because this procedure is invasive and may result in complications¹³. Herein we provide a comprehensive overview of our current understanding of the microflora present in healthy and malignant esophageal tissues, along with insight into other prevalent esophagus-related disorders.

Factors influencing the esophageal microbiome

The esophagus serves as a crucial conduit linking the oral cavity to the stomach, and the esophageal microenvironment undergoes dynamic fluctuations¹⁴. The composition of microbial communities within the esophagus can be influenced by various factors, such as dietary habits. Urban populations, which are characterized by a high consumption of fatty and processed foods, have elevated levels of *Bacteroides*, along with diminished levels of *Firmicutes*. Rural populations adhering to a well-balanced diet rich in fiber have increased levels of *Prevotella*, *Treponema*, and *Succinobacterium*, which aid in the breakdown of polysaccharides and dietary fiber. Prolonged adherence to an urban-style diet can lead to chronic esophageal inflammation and dysbiosis, potentially contributing to the progression of esophageal diseases¹⁵.

Gastroesophageal reflux disease (GERD), a significant risk factor for Barrett's esophagus (BE), can reshape the microecology of the cardia and esophagus due to the reflux content, which is primarily composed of gastric acid and thus creates an acidic environment¹⁴. The impact of medications on the esophageal microbiota should not be underestimated. Proton-pump inhibitors (PPIs) have been shown to cause a notable increase in *Streptococcus* and a decrease in Gram-negative bacteria after use, resulting in reduced inflammation and ulceration in the lower esophagus¹⁶. The combination of omeprazole and antibiotics significantly reduces flora in the lower esophagus of mouse models, leading to the absence of specific bacterial colonization¹⁷. This finding is likely due to the antibiotics targeting *Helicobacter pylori* and other bacteria, or altering the microenvironment in a manner unfavorable for some bacterial populations. Other factors that influence the microbial compositions in the esophagus include obesity, autoimmune disorders, and surgical interventions¹⁸.

Bacterial community of a healthy esophagus

The esophagus, a muscular tube connecting the pharynx and the stomach, serves as a conduit for the transport of food and liquids. The esophagus can be divided into distinct upper, middle, and lower segments. The inner lining of the esophagus is comprised of a stratified squamous epithelium. Investigating

the microbial composition in the esophagus and unraveling the biological functions poses challenges due to its unique anatomy. Traditional culture-based studies have suggested that the esophagus lacks permanent microbial inhabitants, with only a limited presence of transient bacteria acquired through swallowing or gastroesophageal reflux¹⁹.

In 2004, Pei et al.²⁰ identified 95 bacterial species from 6 phyla (*Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria*, *Fusobacteria*, and *TM7*) in the distal esophagus using 16S rDNA sequencing (Figure 1). In 2009, Yang et al.²¹ investigated the distal esophageal microbiota in 12 healthy individuals and

classified the microbiota into 2 subtypes. Type I microbiota is primarily dominated by Gram-positive bacteria, particularly *Streptococcus*, which is typically distributed in the esophagus²¹. Type II microbiome is mainly Gram-negative bacteria presenting in diseased esophagus. *Streptococcus*, along with other genera, such as *Prevotella* and *Lactobacillus*, appears to be one of the dominant taxons in the normal esophageal microbiota, while *Peptostreptococcus*, *Neisseria*, and *Actinobacillus* are less abundant and rarely reported²². In 2013, Norder et al.²³ compared the flora of the lower esophagus, the upper esophagus, and the oral mucosa in healthy individuals, and reported that




Figure 1 Microbiota changes in esophageal cancer. The microbiota of the distal esophagus was influenced by acid reflux from the stomach. Acid reflux leads to inflammation and mucosal damage, resulting in the change of microbiome in the distal esophagus. This process allows the columnar epithelium to replace the original squamous epithelium of the esophagus, which can then progress to BE and EAC. The microbiota in the upper part of the esophagus is influenced by oral resident flora, in which *P. gingivalis* promotes the development of ESCC. BE, Barrett's esophagus; EAC, esophageal adenocarcinoma; EC, esophageal cancer; ESCC, esophageal squamous cell carcinoma.

Table 2 Studies on the esophageal microbiota in healthy individuals

Author, year	Country	Studied subjective (n)	Sampled esophageal tract	Method	Main findings
Gagliardi et al. 1998 ¹⁹	Brazil	Normal esophageal histology (n = 30)	Middle one-third	Culture-based	Esophageal microbiome are acquired through swallowing or gastroesophageal reflux
Pei et al. 2004 ²⁰	US	Normal esophageal histology (n = 4)	2 cm above the squamocolumnar junction	16S rDNA	There are 95 bacterial species from 6 phyla in the distal esophagus
Yang et al. 2009 ²¹	US	Normal (n = 12), esophagitis (n = 12), BE (n = 10)	Distal esophagus	16S rDNA	Esophageal microbiome can be classified into two types (type I and II)
Norder Grusell et al. 2013 ²³	Sweden	Normal esophageal histology (n = 40)	The oral cavity, and upper and lower esophagus	16S rDNA	There are striking similarities in bacterial number and diversity between the oral cavity, and upper and lower esophagus
Dong et al. 2018 ²⁶	China	Normal esophageal histology (n = 27)	The oral cavity, and upper, middle, and lower esophagus	16S rDNA	The microbial composition overlaps between the mouth and esophagus

the composition of the microbiota in these locations was almost comparable. According to a review by Wang et al.,²⁴ other studies have shown that the esophageal microbiota could be affected by adjacent compartments. It is estimated that approximately one bacterial cell per day flows from the mouth to the stomach²⁵, and the microbial composition overlaps between the mouth, pharynx, esophagus, and gut^{14,23}. Specifically, *Streptococcus*, *Neisseria*, *Prevotella*, *Actinomyces*, and *Welchella* are the most abundant species in the oral cavity and esophagus²⁶. The studies that have investigated the microbiota in a healthy esophagus are summarized in **Table 2**.

Esophageal microbial alterations in BE and EAC

Acid reflux-induced inflammation and mucosal damage can lead to GERD and subsequent replacement of the original esophageal squamous epithelium with columnar epithelium that has the potential to progress to BE²⁷ (**Figure 1**). Several studies have identified *Campylobacter* enrichment in BE patients, but not in healthy individuals^{16,21,28}. Conversely, *Siphonobacter*, *Balneola*, *Nitrosopumilus*, and *Planctomyces* have been shown to more abundant in healthy individuals than BE patients²⁹. Yang et al.²¹ demonstrated that BE is primarily associated with an increase in Gram-negative anaerobic bacteria within the esophagus. The presence of lipopolysaccharide (LPS) on the surface of Gram-negative bacteria activates the NF- κ B pathway, leading to elevated expression of IL-8, which is a significant event in the transition from a normal esophagus to BE. Taken together,

these studies suggest that BE-induced microbiome changes in the esophagus may contribute to the development of EAC.

Different studies regarding this hypothesis have generated inconsistent conclusions. A study that involved BE and EAC patients in different stages of disease did not demonstrate a significant difference in the alpha diversities¹³. Snider et al.¹⁷ reported a notable decrease in alpha diversity in high-grade intraepithelial neoplasia and adenocarcinoma compared to non-cancerous esophageal tissue. In addition, a decrease in *Firmicutes* and an increase in *Proteobacteri* were also demonstrated during the progression from BE to EAC¹⁷ (**Figure 1**). Snider et al.¹⁷ also examined changes in the oral microbiota of BE patients, uncovering distinct taxonomic differences, such as an increased abundance of *Streptococcus*, *Veillonella*, and *Enterobacteriaceae*, and decreased *Neisseria*, *Lautropia*, and *Corynebacterium*. Notably, the combination of *Lautropia*, *Streptococcus*, and *Bacteroides* exhibit a high accuracy in identifying BE patients, with a sensitivity of 96.9% and a specificity of 88.2%³⁰. Therefore, alterations in microbiota may serve as a useful tool for diagnosing and monitoring disease progression in patients with BE.

Microbial alterations in EAC

Microbial differences between BE and EAC

BE significantly increases the risk of developing EAC up to 30-fold compared to individuals without BE³¹. Numerous studies have demonstrated notable differences in the microbial composition between healthy individuals, BE patients, and

Table 3 Studies involving the esophageal microbiota in different esophageal diseases

Subtype	Author, year	Country	Sample size	Samples	Method	Alpha diversity	Differentially abundant taxa
BE, GERD and EAC	Yang et al. 2009 ²¹	US	C = 12, ES = 12, BE = 10	Biopsy	16S rDNA	NA	Type II microbiome, such as Gram-negative anaerobes and <i>Streptococcus</i> ↑
	Blackett et al. 2013 ²⁸	US	C = 39, GERD = 37, BE = 45, EAC = 30	Biopsy	16S rDNA	NA	<i>Campylobacteris</i> ↑, <i>H. pylori</i> ↓
	Snider et al. 2019 ¹⁷	US	C = 16, BE = 14, LGD = 6, HGD = 5, EAC = 4	Biopsy	16S rDNA	Lower in EAC	<i>Firmicutes</i> ↑, <i>Proteobacteria</i> ↓
	Elliott et al. 2017 ¹³	UK	C = 20, BE = 24, HGD = 23, EAC = 19	Cyto-sponge, biopsy	16S rDNA	Lower in EAC	<i>Lactobacillus fermentum</i> , <i>Streptococcus</i> ↑
	Peter et al. 2020 ²⁹	US	C = 12, IM = 9, LGD = 12, HGD = 10, EAC = 10	Biopsy, gastric secretions	16S rDNA	NS	<i>Siphonobacter</i> , <i>Balneola</i> , <i>Nitrosopumilus</i> , and <i>Planctomyces</i> ↓
ESCC	Shao et al. 2019 ³⁴	China	ESCC = 45, GCA = 25	Biopsy	16S rDNA	Lower in ESCC	<i>Fusobacterium</i> ↑, <i>Streptococcus</i> ↓
	Li et al. 2020 ³⁵	China	C = 70, ES = 70, LGN = 70, HGN = 19, ESCC = 7	Swab specimens, biopsy	16S rDNA	NS	<i>Streptococcus</i> ↑, <i>Neisseria</i> , and <i>Porphyromonas</i> ↓
	Jiang et al. 2021 ³⁶	China	C = 21, ES = 15, ESCC = 32	Surgical resection, biopsy	16S rDNA	Lower in ESCC	<i>Streptococcus</i> , <i>Actinobacillus</i> , <i>Peptostreptococcus</i> , <i>Fusobacterium</i> , and <i>Prevotella</i> ↑, <i>Faecalibacterium</i> , <i>Bacteroides</i> , <i>Curvibacter</i> and <i>Blautia</i> ↓

C, healthy control; ES, esophagitis; BE, Barrett's esophagus; GERD, gastroesophageal reflux disease; LGD, low grade dysplasia; HGD, high-grade dysplasia; IM, intestinal metaplasia; NA, not available; NS, not significant.

patients with EAC. A comparison of the esophageal microbiota between EAC and BE patients showed reduced diversity in EAC patients, with decreased abundances of *Veillonella* and *Streptococcus granulosa*, while *Lactobacillus* emerged as the dominant flora influencing the local microenvironment¹³ (Figure 1). Of note, the researchers did not detect a significant difference between samples collected from fresh frozen tissues/endoscopic brushings and samples collected using a cyto-sponge, indicating that the cyto-sponge may serve as an alternative method for collecting esophageal microbiota¹³.

Lactobacillus is typically considered to be a part of the resident flora in the stomach. BE originates from the gastric cardia and chronic inflammation in this region exposes *Lactobacillus* to a more acidic environment³². Adaptation of lactic acid bacteria to this low pH environment allows the bacteria to thrive, leading to proliferation and the production of lactic acid through carbohydrate fermentation. This process further acidifies the environment, thus inhibiting the growth of other microorganisms and establishing lactic acid bacteria as the dominant flora³³. The relative abundance of Gram-negative

bacteria and *Enterobacteria* increases simultaneously, progressively intensifying as the disease worsens^{17,29}. Conversely, several bacterial species, such as *Siphonobacter*, *Algae*, *Nitrobacter*, and *Planckia*, are significantly reduced in patients with EAC²⁹ (Figure 1). These findings support the previous study by Yang et al.²¹, which revealed a higher proportion of type II microorganisms in diseased esophagus compared to type I microorganisms. These studies are summarized in Table 3.

Role of *H. pylori* in EAC development

H. pylori has been established as a carcinogen that is closely associated with the progression of various gastric disorders, including gastritis, gastric ulcers, atrophy, and adenocarcinoma³⁷. Although *H. pylori* primarily colonizes the gastric mucosa, the presence of *H. pylori* can influence the microbial composition of the lower esophagus. Tian et al.³⁸ demonstrated that *H. pylori* does not replicate in the esophagus, but has the ability to influence the diversity of the esophageal microbiota. Furthermore, several studies have reported a

lower incidence of EAC in individuals infected with *H. pylori* compared to individuals who are uninfected³⁹⁻⁴¹ (**Figure 1**). This epidemiologic evidence suggests that *H. pylori* infection might be a protective factor against the development of EAC.

Although the precise mechanisms underlying this inverse correlation between *H. pylori* infection and EAC remain unclear, researchers have proposed various explanations for this phenomenon. First, *H. pylori* may counteract the effect of factors that contribute to chronic inflammation and cancer, such as cytotoxin-associated gene A, vacuolated cytotoxin (VAC), and adhesins³⁷. In addition, *H. pylori* has been observed to stimulate cancer cell apoptosis through Fas caspase, thus offering host protection⁴². Third, the presence of *H. pylori* has been linked to serum ghrelin levels. Eradicating *H. pylori* can lead to an increase in serum ghrelin levels, potentially contributing to obesity and affecting gastric emptying, thereby elevating the risk of BE and EAC⁴¹. Furthermore, some studies have linked this inverse correlation with *H. pylori*-induced atrophic gastritis, which can reduce gastric acid secretion and lower the risk of EAC, although this viewpoint remains controversial⁴³.

Microbial alterations in ESCC

Differences between tumor and non-tumor tissues

Shao et al.³⁴ reported reduced microbial diversity in ESCC tissues compared to non-tumor tissues, as determined by 16S rDNA sequencing, with a significant increase in the abundance of *Fusobacterium* and a decrease in *Streptococcus* abundance (**Figure 1**). In 2020, Li et al.³⁵ collected paired saliva and brush specimens from 82 healthy individuals, 60 patients with low-grade dysplasia (LGD), 64 patients with high-grade dysplasia (HGD), and 70 ESCC patients, to examine the microbiota at different stages of ESCC. Li et al.³⁵ revealed significantly decreased abundance of *Streptococcus* and increased abundance of *Neisseria* and *Porphyromonas* during the progression of ESCC. Specifically, *Streptococcus* and *Neisseria* could better predict the development of disease than other genera with reasonable specificity and sensitivity³⁵. Another study conducted by Jiang et al.³⁶ in 2021 that included 68 individuals (controls, $n = 21$; esophagitis, $n = 15$; and ESCC, $n = 32$) who underwent esophagectomy reported contrasting results. Jiang et al.³⁶ reported an increase in *Streptococcus spp.* but a decrease in *Faecalibacterium*, *Bacteroides*, *Curvibacter*, and *Blautia* in ESCC tissues (**Figure 1**). These discrepancies could

be attributed to differences in dietary habits, geographic locations, and variations in the number of patients included in the respective studies. Nevertheless, all these studies collectively demonstrated that alterations in the microbial equilibrium within the esophagus are prevalent in patients with ESCC.

Subsequent studies have indicated that a diminished microbial population serves as a microbial dystrophy index, enhancing the differentiation between EC and a healthy esophagus⁴⁴. Functional analysis of the microbial composition in ESCC has revealed a decline in the activity of nitrate and nitrite reductases, which are closely linked to carcinogenesis through the production of reactive nitrates and nitrites⁴⁵. Although these findings imply that microbial dysbiosis may contribute to the development of ESCC, further research is warranted to ascertain the specific microbiota involved and elucidate the underlying mechanisms.

Influence of *Porphyromonas gingivalis* in ESCC development

In 2016, Gao et al.⁴⁶ investigated the presence of antigens, DNA, and periodontal pathogens in ESCC lesions and showed that the presence of *P. gingivalis* is more prevalent in ESCC than para-cancerous tissues (61% vs. 12%) but absent in the esophageal epithelium⁴⁶ (**Figure 1**). Furthermore, the abundance of *P. gingivalis* varies across different stages of ESCC, with higher levels observed in patients with poor differentiation, severe lymph node metastasis, advanced stage disease, and a short survival cycle. These findings suggested that *P. gingivalis* could serve as a novel prognostic indicator for ESCC. Peters et al.⁴⁷ determined the oral bacteria present in 25 pairs of patients with ESCC and healthy controls. The study revealed a higher prevalence of *P. gingivalis* in ESCC tumor tissues compared to paired healthy controls⁴⁷. In addition, elevated levels of IgG and IgA antibodies against *P. gingivalis* were detected in ESCC patients compared to healthy controls. Notably, patients with high antibody levels exhibited a more favorable prognosis than patients with low antibody levels. These findings suggested that *P. gingivalis* may have a role in the pathogenesis and progression of ESCC⁴⁸.

A recent study performed by Chen et al.⁴⁹ determined the presence of *P. gingivalis* in the esophagus of 156 ESCC patients using immunohistochemistry to investigate the possible association between *P. gingivalis* infection and patient clinicopathologic features. Chen et al.⁴⁹ detected *P. gingivalis* in 57% of the ESCC patients, and the infection contributed to EC development by promoting IL-6 production to induce

the epithelial-mesenchymal transition and attract myeloid-derived suppressor cells. It is worth noting that *P. gingivalis* interacts with EC epithelial cells in different ways. For example, *P. gingivalis* secretes nucleoside diphosphate kinase to promote carcinogenesis⁵⁰ and inhibits epithelial cell apoptosis through different pathways, such as activation of Jak1/Akt/Stat3 signaling⁵¹, enhancement of Bcl-2, and blocking the release of cytochrome c⁵² (Figure 1). Therefore, it is reasonable to consider *P. gingivalis* as a promising target to prevent and/or treat *P. gingivalis*-infected patients with ESCC⁴⁹.

Microbiota in association with the prognosis of ESCC

Liu et al.⁵³ explored the presence of esophageal microbiota in ESCC patients at different pathologic stages in an attempt to identify potential microbial markers with prognostic value. The findings revealed significant differences in the abundance of bacterial phyla and genera between patients with lymph node metastasis (N+) and patients without lymph node metastasis (N-). Notably, *Bacteroidetes*, *Pleurotus*, and *Spirochetes* had a higher abundance in N+ patients, while *Proteobacteria* exhibited a lower abundance in N+ patients compared to N- patients (Figure 1). At the genus level, *Prevotella* and *Treponema* were more abundant in the N+ group, while *Streptococcus* exhibited a higher abundance in patients with T3-4 tumors compared to T1-2 tumors. No significant differences were observed in the abundance of other genera. Additionally, the analysis indicated that the combined abundance of *Streptococcus* and *Prevotella* was associated with poor survival, suggesting that these genera could potentially serve as independent prognostic indicators for ESCC⁵³.

Periodontal disease has been recognized as a significant risk factor for EC. Recently, increasing attention has been given to the potential involvement of oral flora in EC development. Notably, *Fusobacterium nucleatum*, a pathogen commonly associated with periodontal disease and colorectal cancer⁵⁴, has been extensively studied (Figure 1). Yamamura et al.⁵⁵ detected *F. nucleatum* DNA using PCR in 23% (74/325) of ESCC resected specimens, primarily from advanced-stage cases. The presence of *F. nucleatum* appeared to be independent of other factors, such as gender, age, cigarette smoking, alcohol consumption, and preoperative treatment⁵⁶. Moreover, higher levels of *F. nucleatum* were observed more often in relapsed cases, and these patients had a lower survival rate compared to *F. nucleatum*-negative individuals⁵⁵. Subsequent investigations

focusing on ESCC patients after radiotherapy and chemotherapy revealed that those with a high abundance of *F. nucleatum* experienced more adverse reactions to chemotherapy, had a higher recurrence rate, and a shorter survival time compared to patients with a low *F. nucleatum* abundance, suggesting the potential contribution to chemoresistance⁵⁷. These findings collectively support the notion that *F. nucleatum* holds promise as a prognostic factor for ESCC.

Effect of human papillomavirus (HPV) infection on ESCC

HPV infection has been closely linked to oropharyngeal squamous cell carcinoma⁵⁸. Considering the similarity in human leukocyte antigen expression between squamous epithelial cells in the oropharynx and esophagus, it is reasonable to propose that HPV can potentially contribute to the development of ESCC. A case-control study conducted in Shaanxi, China aimed to detect HPV antigens in the blood and revealed significantly higher serum reactivity in ESCC patients compared to healthy controls⁵⁹; however, subsequent studies on this topic have yielded conflicting results. Specifically, a study by Kamangar et al.³⁹ in 2006 detected HPV antibodies in < 15% of serum samples from Chinese ESCC patients, and no definitive correlation between HPV and ESCC was established. Similarly, a study by Halec et al.⁶⁰ in 2016 investigating the potential association between HPV and ESCC found no significant association. These studies used different methods to detect HPV infection in ESCC tissues and consistently observed low viral loads in cancer tissues. Furthermore, the presence of HPV DNA, mRNA, and subsequent p16 upregulation was not consistently observed. Based on the collective findings, the association between HPV infection and the incidence of ESCC appears to be weak.

In contrast, Sitas et al.⁶¹ utilized a centralized multiple serology method to analyze sera from 1561 patients with ESCC and 2502 controls. Sitas et al.⁶¹ found that ESCC was only associated with E6 for HPV16 and HPV6, but not with other types of HPV. Similarly, a study by Zhang et al.⁶² concluded that only individuals who were cigarette smokers, consumed alcohol, and were infected with HPV had a higher likelihood of developing ESCC. This finding suggested that HPV infection alone may not be an independent risk factor for ESCC but could potentially have a synergistic effect with other factors. The current epidemiologic and etiologic evidence supporting the association between HPV infection and ESCC remains inconclusive, and the topic continues to be a subject of discussion.

Dysbiosis facilitates the development of EC by disturbing the immune response

Changes in the composition and abundance of esophageal microbiota could promote the development of EC in different ways. For example, several lactic acid-producing bacteria, such as *Staphylococcus* and *Lactobacillus*, have an increased

abundance in EAC tissues, which converts the high load of glucose taken up by cancer cells into lactate, thereby supporting the survival and proliferation of malignant cells⁶³ (Figure 2). While extensive research is currently focused on investigating the precise mechanisms by which esophageal microbiota contribute to the development of EC, a prevailing observation is that the presence of chronic inflammation and compromised immune responses establishes a conducive microenvironment for this malignant transformation. The involvement




Figure 2 A diagram on the interactions between esophageal microbiota and immune cells to promote the development of EC. Changes in the composition and abundance of esophageal microbiota promote the development of EC in different ways. Notably, increased lactic acid-producing bacteria, such as *Staphylococcus* and *Lactobacillus* in EC tissues, support tumor survival and proliferation by converting glucose into lactate. Chronic inflammation and compromised immune responses in the esophageal microenvironment create a conducive setting for malignant transformation. Dysregulation of the Wnt/ β -catenin pathway by *Fusobacterium nucleatum* contributes to disease progression and treatment resistance through elevated production of chemokines. The altered esophageal microbiota activates multiple TLRs (TLR1, 2, and 6) and NLRP3 (a component of inflammasomes) to disturb the local microenvironment homeostasis and promote malignant cellular behavior.

of disrupted cross-talk between esophageal microbiota and immune cells has been implicated in several signaling pathways known to contribute to the development of EC.

Aberrant activation of the Wnt/ β -catenin pathway has been implicated in the carcinogenesis and therapeutic resistance of EC⁶⁴. *Fusobacterium nucleatum*, a bacterium capable of activating the β -catenin pathway through the production of FadA adhesion and modulating the inflammatory response, has been reported to be present in approximately 23% of patients with EC⁵⁵. The tumor tissues of these affected patients have increased production of chemokines, which contributes to a more aggressive disease course and reduced survival⁵⁵. In addition, changes in esophageal microbiota have been shown to facilitate the activation of multiple Toll-like receptors (TLRs) through different approaches. TLR4 activation led to increased NF- κ B activities and cyclooxygenase-2 (COX-2) expression. The former promotes secretion of several chemokines, such as IL-6, IL-8, and TNF, while the latter is known to be associated with different malignant cellular phenotypes, including increased proliferation, angiogenesis, invasiveness, and decreased apoptosis⁶⁵. TLR1, 2, and 6 induce inflammatory responses, the upregulation of which has been reported in EAC tissues. Of note, the TLR1, 2, and 6 network has also been implicated in identifying dysbiotic microbial components⁶⁶. Third, Gram-negative bacteria stimulate inducible nitric oxide synthase (iNOS), the expression of which has been shown to be increased in EAC tissues compared to normal esophagus⁶⁷. Depletion of Nod-like receptor protein 3 (NLRP3), a component of inflammasomes, causes aberrant bactericidal activities. Conversely, NLRP3 activation allows Tregs to maintain homeostasis by enhancing the secretion of IL-1 β to neutralize the inflammatory response⁶⁸ (Figure 2). Taken together, these studies indicate a close relationship between esophageal dysbiosis and aberrant immune responses, ultimately leading to the malignant transformation of EC, as reviewed by Sharma et al.⁶⁹

Summary and outlook

The field of esophageal microbiology remains relatively unexplored, and the causal relationship between EC and the microbiome remains uncertain. Nevertheless, the studies reviewed in this paper have provided insight into the microbial composition of the normal esophagus and identified changes in microbial composition among EC patients, which is largely characterized by a reduction in bacterial species. There is no

doubt that emerging advances in technology and innovative tools are progressively enhancing our understanding of the intricate involvement of the microbiota in the pathogenesis of EC. Cyto-sponge, as a minimally invasive method for microbiota sampling, demonstrated superior performance by yielding higher quantities of microbial DNA and capturing a comparable microbial profile to biopsy and brush samples, while exhibiting an enrichment of taxa from the oral cavity and stomach¹³. In addition, organoids have emerged as a valuable *ex vivo* tool for modeling esophageal homeostasis and disease, faithfully reproducing the dynamic characteristics of the esophageal epithelium. Organoids successfully recapitulate normal epithelial renewal, differentiation, and proliferation, making organoids suitable for studying disease-specific alterations in response to various pathogenic stimuli⁷⁰. Co-culture models of 3D organoids and the gut microbiome have enabled the faithful characterization of the consequences of microbe-epithelial interactions⁷¹. It is clear that 3D organoid models are an ideal platform for examining host-pathogen interactions in the co-culture of the microbiome and esophageal tissue. In addition, the escalating adoption of target therapies in EC has led to accumulating evidence suggesting the potential involvement of esophageal microbiota in modulating the patient response to these treatments^{72,73}, thus presenting a significant focus of research in this field.

Our exploration of the association between the microbiome and EC has considered various factors that can influence changes in the esophageal microbial community, including medications, immune response, dietary habits, and age. Additionally, we have investigated the oral microbiome, which is suspected to be a risk factor for EC, particularly ESCC. However, there are discrepancies in these findings, and significant gaps persist in our understanding of the etiology, pathology, and immunology of this disease. Therefore, large-scale prospective cohort studies are invaluable to monitor the longitudinal changes in the microbiome during lesion progression and intervention. The current sampling methods for esophageal flora are becoming more diverse, and the use of non-endoscopic cell sampling device, like the cyto-sponge, offers non-invasive and convenient sampling options that can facilitate further exploration of the esophageal flora.

Grant support

This work was supported by grants from the Health Commission of Henan Province (Grant No. SBGJ20211008)

and the Henan Provincial Department of Science and Technology (Grant No. 222300420574).

Conflict of interest statement

No potential conflicts of interest are disclosed.

Author contributions

Conceived and designed the analysis: Jun Li, Hongle Li.

Collected the data: Yi Li.

Contributed data or analysis tools: Xia Xue, Yi Li, Bing Wei.

Performed the analysis: Yi Li, Jun Li.

Wrote the paper: Yi Li, Jun Li.

References

- Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global cancer statistics 2020: Globocan estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin.* 2021; 71: 209-49.
- Abnet CC, Arnold M, Wei WQ. Epidemiology of esophageal squamous cell carcinoma. *Gastroenterology.* 2018; 154: 360-73.
- Uhlenhopp DJ, Then EO, Sunkara T, Gaduputi V. Epidemiology of esophageal cancer: update in global trends, etiology and risk factors. *Clin J Gastroenterol.* 2020; 13: 1010-21.
- Rubenstein JH, Shaheen NJ. Epidemiology, diagnosis, and management of esophageal adenocarcinoma. *Gastroenterology.* 2015; 149: 302-17.e1.
- Shapiro J, van Lanschot JJB, Hulshof M, van Hagen P, van Berge Henegouwen MI, Wijnhoven BPL, et al. Neoadjuvant chemoradiotherapy plus surgery versus surgery alone for oesophageal or junctional cancer (CROSS): long-term results of a randomised controlled trial. *Lancet Oncol.* 2015; 16: 1090-8.
- Zhang Y. Epidemiology of esophageal cancer. *World J Gastroenterol.* 2013; 19: 5598-606.
- Wheeler JB, Reed CE. Epidemiology of esophageal cancer. *Surg Clin North Am.* 2012; 92: 1077-87.
- Domper Arnal MJ, Ferrández Arenas Á, Lanás Arbeloa Á. Esophageal cancer: risk factors, screening and endoscopic treatment in western and eastern countries. *World J Gastroenterol.* 2015; 21: 7933-43.
- Wei WQ, Abnet CC, Lu N, Roth MJ, Wang GQ, Dye BA, et al. Risk factors for oesophageal squamous dysplasia in adult inhabitants of a high risk region of china. *Gut.* 2005; 54: 759-63.
- Schneider JL, Corley DA. A review of the epidemiology of Barrett's oesophagus and oesophageal adenocarcinoma. *Best Pract Res Clin Gastroenterol.* 2015; 29: 29-39.
- Gilbert JA, Quinn RA, Debelius J, Xu ZZ, Morton J, Garg N, et al. Microbiome-wide association studies link dynamic microbial consortia to disease. *Nature.* 2016; 535: 94-103.
- Zou S, Fang L, Lee MH. Dysbiosis of gut microbiota in promoting the development of colorectal cancer. *Gastroenterol Rep.* 2018; 6: 1-12.
- Elliott DRE, Walker AW, O'Donovan M, Parkhill J, Fitzgerald RC. A non-endoscopic device to sample the oesophageal microbiota: a case-control study. *Lancet Gastroenterol Hepatol.* 2017; 2: 32-42.
- Gall A, Fero J, McCoy C, Claywell BC, Sanchez CA, Blount PL, et al. Bacterial composition of the human upper gastrointestinal tract microbiome is dynamic and associated with genomic instability in a Barrett's esophagus cohort. *PLoS ONE.* 2015; 10: e0129055.
- Nobel YR, Snider EJ, Compres G, Freedberg DE, Khiabani H, Lightdale CJ, et al. Increasing dietary fiber intake is associated with a distinct esophageal microbiome. *Clin Transl Gastroenterol.* 2018; 9: 199.
- Amir I, Konikoff FM, Oppenheim M, Gophna U, Half EE. Gastric microbiota is altered in oesophagitis and Barrett's oesophagus and further modified by proton pump inhibitors. *Environ Microbiol.* 2014; 16: 2905-14.
- Snider EJ, Compres G, Freedberg DE, Khiabani H, Nobel YR, Stump S, et al. Alterations to the esophageal microbiome associated with progression from Barrett's esophagus to esophageal adenocarcinoma. *Cancer Epidemiol Biomarkers Prev.* 2019; 28: 1687-93.
- Dominguez-Bello MG, Godoy-Vitorino F, Knight R, Blaser MJ. Role of the microbiome in human development. *Gut.* 2019; 68: 1108-14.
- Gagliardi D, Makihara S, Corsi PR, Viana Ade T, Wiczler MV, Nakakubo S, et al. Microbial flora of the normal esophagus. *Dis Esophagus.* 1998; 11: 248-50.
- Pei Z, Bini EJ, Yang L, Zhou M, Francois F, Blaser MJ. Bacterial biota in the human distal esophagus. *Proc Natl Acad Sci U S A.* 2004; 101: 4250-5.
- Yang L, Lu X, Nossa CW, Francois F, Peek RM, Pei Z. Inflammation and intestinal metaplasia of the distal esophagus are associated with alterations in the microbiome. *Gastroenterology.* 2009; 137: 588-97.
- Di Pilato V, Freschi G, Ringressi MN, Pallecchi L, Rossolini GM, Bechi P. The esophageal microbiota in health and disease. *Ann NY Acad Sci.* 2016; 1381: 21-33.
- Norder Grusell E, Dahlén G, Ruth M, Ny L, Quiding-Järbrink M, Bergquist H, et al. Bacterial flora of the human oral cavity, and the upper and lower esophagus. *Dis Esophagus.* 2013; 26: 84-90.
- Wang WL, Xu SY, Ren ZG, Tao L, Jiang JW, Zheng SS. Application of metagenomics in the human gut microbiome. *World J Gastroenterol.* 2015; 21: 803-14.
- Segata N, Haake SK, Mannon P, Lemon KP, Waldron L, Gevers D, et al. Composition of the adult digestive tract bacterial microbiome based on seven mouth surfaces, tonsils, throat and stool samples. *Genome Biol.* 2012; 13: R42.
- Dong L, Yin J, Zhao J, Ma SR, Wang HR, Wang M, et al. Microbial similarity and preference for specific sites in healthy oral cavity and esophagus. *Front Microbiol.* 2018; 9: 1603.
- Liu N, Ando T, Ishiguro K, Maeda O, Watanabe O, Funasaka K, et al. Characterization of bacterial biota in the distal esophagus of

- Japanese patients with reflux esophagitis and Barrett's esophagus. *BMC Infect Dis.* 2013; 13: 130.
28. Blackett KL, Siddhi SS, Cleary S, Steed H, Miller MH, Macfarlane S, et al. Oesophageal bacterial biofilm changes in gastro-oesophageal reflux disease, Barrett's and oesophageal carcinoma: association or causality? *Aliment Pharmacol Ther.* 2013; 37: 1084-92.
 29. Peter S, Pendergraft A, VanDerPol W, Wilcox CM, Kyanam Kabir Baig KR, Morrow C, et al. Mucosa-associated microbiota in Barrett's esophagus, dysplasia, and esophageal adenocarcinoma differ similarly compared with healthy controls. *Clin Transl Gastroenterol.* 2020; 11: e00199.
 30. Snider EJ, Compres G, Freedberg DE, Giddins MJ, Khiabani H, Lightdale CJ, et al. Barrett's esophagus is associated with a distinct oral microbiome. *Clin Transl Gastroenterol.* 2018; 9: 135.
 31. Cook MB, Coburn SB, Lam JR, Taylor PR, Schneider JL, Corley DA. Cancer incidence and mortality risks in a large US Barrett's oesophagus cohort. *Gut.* 2018; 67: 418-529.
 32. McDonald SA, Lavery D, Wright NA, Jansen M. Barrett oesophagus: lessons on its origins from the lesion itself. *Nat Rev Gastroenterol Hepatol.* 2015; 12: 50-60.
 33. Kleerebezem M, Hols P, Bernard E, Rolain T, Zhou M, Siezen RJ, et al. The extracellular biology of the lactobacilli. *FEMS Microbiol Rev.* 2010; 34: 199-230.
 34. Shao D, Vogtmann E, Liu A, Qin J, Chen W, Abnet CC, et al. Microbial characterization of esophageal squamous cell carcinoma and gastric cardia adenocarcinoma from a high-risk region of China. *Cancer.* 2019; 125: 3993-4002.
 35. Li M, Shao D, Zhou J, Gu J, Qin J, Chen W, et al. Signatures within esophageal microbiota with progression of esophageal squamous cell carcinoma. *Chin J Cancer Res.* 2020; 32: 755-67.
 36. Jiang Z, Wang J, Shen Z, Zhang Z, Wang S. Characterization of esophageal microbiota in patients with esophagitis and esophageal squamous cell carcinoma. *Front Cell Infect Microbiol.* 2021; 11: 774330.
 37. Dzutsev A, Goldszmid RS, Viaud S, Zitvogel L, Trinchieri G. The role of the microbiota in inflammation, carcinogenesis, and cancer therapy. *Eur J Immunol.* 2015; 45: 17-31.
 38. Tian Z, Yang Z, Gao J, Zhu L, Jiang R, Jiang Y. Lower esophageal microbiota species are affected by the eradication of helicobacter pylori infection using antibiotics. *Exp Ther Med.* 2015; 9: 685-92.
 39. Kamangar F, Qiao YL, Schiller JT, Dawsey SM, Fears T, Sun XD, et al. Human papillomavirus serology and the risk of esophageal and gastric cancers: results from a cohort in a high-risk region in China. *Int J Cancer.* 2006; 119: 579-84.
 40. Anderson LA, Murphy SJ, Johnston BT, Watson RG, Ferguson HR, Bamford KB, et al. Relationship between Helicobacter pylori infection and gastric atrophy and the stages of the oesophageal inflammation, metaplasia, adenocarcinoma sequence: results from the FINBAR case-control study. *Gut.* 2008; 57: 734-9.
 41. Nie S, Chen T, Yang X, Huai P, Lu M. Association of helicobacter pylori infection with esophageal adenocarcinoma and squamous cell carcinoma: a meta-analysis. *Dis Esophagus.* 2014; 27: 645-53.
 42. Tözün N, Vardareli E. Gut microbiome and gastrointestinal cancer: Les Liaisons Dangereuses. *J Clin Gastroenterol.* 2016; 50 Suppl 2.
 43. Bik EM, Eckburg PB, Gill SR, Nelson KE, Purdom EA, Francois F, et al. Molecular analysis of the bacterial microbiota in the human stomach. *Proc Natl Acad Sci U S A.* 2006; 103: 732-7.
 44. Yang W, Chen CH, Jia M, Xing X, Gao L, Tsai HT, et al. Tumor-associated microbiota in esophageal squamous cell carcinoma. *Front Cell Dev Biol.* 2021; 9: 641270.
 45. Lin Z, Rao W, Xiang Z, Zeng Q, Liu S, Yu K, et al. Characteristics and interplay of esophageal microbiota in esophageal squamous cell carcinoma. *BMC Cancer.* 2022; 22: 696.
 46. Gao S, Li S, Ma Z, Liang S, Shan T, Zhang M, et al. Presence of porphyromonas gingivalis in esophagus and its association with the clinicopathological characteristics and survival in patients with esophageal cancer. *Infect Agent Cancer.* 2016; 11: 3.
 47. Peters BA, Wu J, Pei Z, Yang L, Purdue MP, Freedman ND, et al. Oral microbiome composition reflects prospective risk for esophageal cancers. *Cancer Res.* 2017; 77: 6777-87.
 48. Gao SG, Yang JQ, Ma ZK, Yuan X, Zhao C, Wang GC, et al. Preoperative serum immunoglobulin G and A antibodies to Porphyromonas gingivalis are potential serum biomarkers for the diagnosis and prognosis of esophageal squamous cell carcinoma. *BMC Cancer.* 2018; 18: 17.
 49. Chen MF, Lu MS, Hsieh CC, Chen WC. Porphyromonas gingivalis promotes tumor progression in esophageal squamous cell carcinoma. *Cell Oncol (Dordrecht).* 2021; 44: 373-84.
 50. Morandini AC, Ramos-Junior ES, Potempa J, Nguyen KA, Oliveira AC, Bellio M, et al. Porphyromonas gingivalis fimbriae dampen P2x7-dependent interleukin-1 β secretion. *J Innate Immun.* 2014; 6: 831-45.
 51. Yilmaz O, Jungas T, Verbeke P, Ojcius DM. Activation of the phosphatidylinositol 3-kinase/Akt pathway contributes to survival of primary epithelial cells infected with the periodontal pathogen Porphyromonas gingivalis. *Infect Immun.* 2004; 72: 3743-51.
 52. Yao L, Jermanus C, Barbetta B, Choi C, Verbeke P, Ojcius DM, et al. Porphyromonas gingivalis infection sequesters pro-apoptotic Bad through Akt in primary gingival epithelial cells. *Mol Oral Microbiol.* 2010; 25: 89-101.
 53. Liu Y, Lin Z, Lin Y, Chen Y, Peng XE, He F, et al. Streptococcus and prevotella are associated with the prognosis of oesophageal squamous cell carcinoma. *J Med Microbiol.* 2018; 67: 1058-68.
 54. Flanagan L, Schmid J, Ebert M, Soucek P, Kunicka T, Liska V, et al. Fusobacterium nucleatum associates with stages of colorectal neoplasia development, colorectal cancer and disease outcome. *Eur J Clin Microbiol Infect Dis.* 2014; 33: 1381-90.
 55. Yamamura K, Baba Y, Nakagawa S, Mima K, Miyake K, Nakamura K, et al. Human microbiome fusobacterium nucleatum in esophageal cancer tissue is associated with prognosis. *Clin Cancer Res.* 2016; 22: 5574-81.
 56. Nomoto D, Baba Y, Liu Y, Tsutsuki H, Okadome K, Harada K, et al. Fusobacterium nucleatum promotes esophageal squamous cell carcinoma progression via the NOD1/RIPK2/NF- κ B pathway. *Cancer Lett.* 2022; 530: 59-67.

57. Yamamura K, Izumi D, Kandimalla R, Sonohara F, Baba Y, Yoshida N, et al. Intratumoral fusobacterium nucleatum levels predict therapeutic response to neoadjuvant chemotherapy in esophageal squamous cell carcinoma. *Clin Cancer Res.* 2019; 25: 6170-9.
 58. Wang WL, Wang YC, Chang CY, Lo JL, Kuo YH, Hwang TZ, et al. Human papillomavirus infection on initiating synchronous esophageal neoplasia in patients with head and neck cancer. *Laryngoscope.* 2016; 126: 1097-102.
 59. Han C, Qiao G, Hubbert NL, Li L, Sun C, Wang Y, et al. Serologic association between human papillomavirus type 16 infection and esophageal cancer in Shaanxi Province, China. *J Natl Cancer Inst.* 1996; 88: 1467-71.
 60. Halec G, Schmitt M, Egger S, Abnet CC, Babb C, Dawsey SM, et al. Mucosal alpha-papillomaviruses are not associated with esophageal squamous cell carcinomas: lack of mechanistic evidence from South Africa, China and Iran and from a world-wide meta-analysis. *Int J Cancer.* 2016; 139: 85-98.
 61. Sitas F, Egger S, Urban MI, Taylor PR, Abnet CC, Boffetta P, et al. InterSCOPE study: associations between esophageal squamous cell carcinoma and human papillomavirus serological markers. *J Natl Cancer Inst.* 2012; 104: 147-58.
 62. Zhang SK, Guo LW, Chen Q, Zhang M, Liu SZ, Quan PL, et al. The association between human papillomavirus 16 and esophageal cancer in Chinese population: a meta-analysis. *BMC Cancer.* 2015; 15: 1096.
 63. Deshpande NP, Riordan SM, Castaño-Rodríguez N, Wilkins MR, Kaakoush NO. Signatures within the esophageal microbiome are associated with host genetics, age, and disease. *Microbiome.* 2018; 6: 227.
 64. Das PK, Islam F, Smith RA, Lam AK. Therapeutic strategies against cancer stem cells in esophageal carcinomas. *Front Oncol.* 2020; 10: 598957.
 65. Abdel-Latif MM, Kelleher D, Reynolds JV. Potential role of NF-kappaB in esophageal adenocarcinoma: as an emerging molecular target. *J Surg Res.* 2009; 153: 172-80.
 66. Verbeek RE, Siersema PD, Vleggaar FP, Ten Kate FJ, Posthuma G, Souza RE, et al. Toll-like receptor 2 signalling and the lysosomal machinery in Barrett's esophagus. *J Gastrointest Liver Dis.* 2016; 25: 273-82.
 67. Clemons NJ, Shannon NB, Abeyratne LR, Walker CE, Saadi A, O'Donovan ML, et al. Nitric oxide-mediated invasion in Barrett's high-grade dysplasia and adenocarcinoma. *Carcinogenesis.* 2010; 31: 1669-75.
 68. Nadatani Y, Huo X, Zhang X, Yu C, Cheng E, Zhang Q, et al. NOD-like receptor protein 3 inflammasome priming and activation in Barrett's epithelial cells. *Cell Mol Gastroenterol Hepatol.* 2016; 2: 439-53.
 69. Sharma T, Gupta A, Chauhan R, Bhat AA, Nisar S, Hashem S, et al. Cross-talk between the microbiome and chronic inflammation in esophageal cancer: potential driver of oncogenesis. *Cancer Metastasis Rev.* 2022; 41: 281-99.
 70. Flashner S, Yan KS, Nakagawa H. 3D organoids: an untapped platform for studying host-microbiome interactions in esophageal cancers. *Microorganisms.* 2021; 9: 2182.
 71. Min S, Kim S, Cho SW. Gastrointestinal tract modeling using organoids engineered with cellular and microbiota niches. *Exp Mol Med.* 2020; 52: 227-37.
 72. Bai M, Wang M, Deng T, Bai Y, Zang K, Miao Z, et al. Safety and efficacy of anti-EGFR monoclonal antibody (SCT200) as second-line therapy in advanced esophageal squamous cell carcinoma. *Cancer Biol Med.* 2022; 19: 358-69.
 73. Yin H, Yang L, Peng G, Yang K, Mi Y, Hu X, et al. The commensal consortium of the gut microbiome is associated with favorable responses to anti-programmed death protein 1 (PD-1) therapy in thoracic neoplasms. *Cancer Biol Med.* 2021; 18: 1040-52.
- Cite this article as:** Li Y, Wei B, Xue X, Li H and Li J. Microbiome changes in esophageal cancer: implications for pathogenesis and prognosis. *Cancer Biol Med.* 2023; xx: xx-xx. doi: 10.20892/j.issn.2095-3941.2023.0177